



IRAQI
Academic Scientific Journals



العراقية
المجلات الأكاديمية العلمية

TJAS
Tikrit Journal for
Agricultural
Sciences

ISSN:1813-1646 (Print); 2664-0597 (Online)

Tikrit Journal for Agricultural Sciences

Journal Homepage: <http://www.tjas.org>

E-mail: tjas@tu.edu.iq

**Mijbil Mohammad
Aljumaily
Hudhaifa Maan
Al-Hamandi ***

*Department of Soil
Science and Water
Resources, College of
Agriculture, Tikrit
University, Tikrit, Iraq*

KEY WORDS:

organic substances,
metal complexes,
Stability constant, FA-
metal complex

Hu-metal complex.

ARTICLE HISTORY:

Received: 16/08/2022

Accepted: 07/09/2022

Available online:
30/9/2022

Tikrit Journal for Agricultural Sciences (TJAS)

Organic Matter and Heavy Metals Sorption

ABSTRACT

Organic matter content in soils is highly variable and includes dead and living organisms and their decomposition products. plant residue and humic substances. Thermodynamically, organic matter is unstable in soils and later will oxidize to CO_2 , and H_2O . The effective substances of organic matter decomposition are fulvic and humic acids (FA+Hu) which contain Several functional groups that release electrons or protons during their decomposition leaving behind several radical groups that act as electron donor to ward heavy metal ions forming FA and Hu-metal soluble and insoluble complexes. Those metallic-organic Complexes are variable by their Stability constant (SC) which is absolutely pH-dependent. The less stable the metal complex, the higher mobility in soil, In contrast, a highly stable metal complex is less soluble and mobility. So, organic matter plays an important role in the accumulation, leaching, and transportation of heavy metal Cations present in water and soils as chelates of different Stability and supplying plant roots by these ions and behavior as a buffering substance to heavy metal mobility.

© 2022 TJAS. College of Agriculture, Tikrit University

INTRODUCTION

Organic matter (OM) is a wide term include residual of plants, animals and microorganisms with different degree of decomposition (Walker et al. 2003; Liu et al. 2022). Soil organic matter (SOM) is a complex substances which simplified easily by soil microorganism to allow molecular weight materials (Bolan and Duraisamy, 2003). In the most soils, the final decomposition product and/or by product of (OM) results are, humic acid organic acids of high and low molecular weights, amino acids, waxes, protein, lipids, peptides, lignin, aromatic and aliphatic hydrocarbons (AL- Hamandi, 2018). The final decomposition of (SOM) can be divided into - fulvic acid (FA), humic acid (HA) and humine which widely exist in soil and presents an important fraction of soil component (Weber, 1988). Earth crust containing about (50×10^6 tan) of organic matter as humus(Killops and Killops,2013). The general characteristic's of humic substances are:- increasing dissolved organic substances. Laboratory practical experiments concluded that humic acid reduces many heavy metal ions such as (Fe^{+3} to Fe^+ , Cu^{+2} to Cu^0 , and Hg^{+2} to Hg^0 (Matthews,1983). While Minkina. et, al (2006) reported that (SOM) interactive, With Zn, Pb, and Cu resulted changes in (SOM) product, reduce (HA) and increased (FA) content.

* Corresponding author: E-mail: hudhaifaalhamandi@tu.edu.iq

Definitions of the components of organic matter

- Fulvic acids (fulvate):- have a high acidity with less degree of polymerization, high mobility, and exist in acid soils with low biological activity
- Humic acids (humates):- Spherocolloide shape, medium mobility, acidity, and occur in neutral soils with high biological activity.
- Humins: have low acidity, and a high degree of polymerization and occur in every soil (Kabata-pendias, 2000) cation exchange capacity (CEC) for humic Substances are vary between 200- 450 C.mole Kg-1, being lower for (FAS) and higher for (HAS).

Soil properties are controlled by soil organic matter (SOM) because these substances increase soil CEC from 20% to 70% (Evans et al, 2003). This great Sorption Capacity for heavy metals is very important in reducing the activity of excess heavy metals. (Laxen, 1985). Cu, Ni Cr, and Cd are enhanced Significantly by heavy metals interaction with humic Substances which have been described as chelation. Coagulation, Surface Sorption, and ion exchange (Tan, 2010). The reactions between cations and organic substances lead to the form of water-soluble and insoluble compounds and some of the organic matter product acts as reducing materials to enhance the mobile Zing of trace elements (McBride et al, 1997) (Table1). Dissolved organic Substances could effectively form complexes with heavy metal ions in Soil (Bolton and Evans, 1991) Which Can be moved through the Soil matrix downward and may contaminate groundwater (Li and Shuman, 1996). Also Boyle and Fuller (1987), Lamy et al. (1993) reported that Zn and cd mobility was enhanced by heavy metal ions tend to accumulate in the products of organic matter decompositions, products with fewer metal ion content, and the higher binding energy between metal cans and organic groups (Zunino et al.1979). In order to be interested and some information about insoluble complete behavior, we should seek by a complex Stability constant (CSC) concept which can be identified as an equilibrium constant that forms a chelate or soluble complex the stability index value has been used by Halder et al (2020). Which describes the ratio of metal-OM to the amount of metal in inorganic fractions. Pure fulvic acid (FA) and humic acid (HA) stability index show that heavy metal ions of Mn^{+2} , Zn^{+2} , Pb^{+2} , and Cu^{+2} are more ready to form complex with (HA) than with (FA) and over pH (4 to5) large quantities of Cu^{+2} are complexed with (HA), but at pH range between (6-7) Cu^{+2} ions complexed with FA. Both heavy metal ions Pb^{+2} and Cu^{+2} showed high affinity to complicated organic acids compared to Mn^{+2} and Fe^{+2} ions (Kabata-pendias, 2000). The findings reported by Förstner and Müller (2013) Pauli (1975) ; Van Dijk (1971). Stevenson (1972), Vlasov and Michylova (1975) indicate that the stability constant of metallo-organic complexes, although variable, pH depending, and other medium properties can be ordered as the following sequence:



Decomposition of O.M results in many organic substances, such as phenols, hydroxy acids, amino acids, and phosphoric acids which naturally exist in soils. These compounds behave as a chelating agent for heavy metals. Metal complex's solubility depends on binding Strength and formed complexes which are determined by the organic group's size and mobility of formed complexes (Kabata-penelias, 2000).

Metallio-organic and pH dependent

Chelation and complexation of heavy metal ions with organic substances in water have been studied by Sholkovitz and Copland (1981). They reported that the solubilities of (HA) complexes with (Cd, Ni, Cu, Mn, and Fe) differ from that concluded from inorganic Solubility consideration. These ions complexing with humic substances are solubilized at (Moderate-high) PH ranging between (3-9) and precipitate at low pH values ranging between (1.5-3.5). Thus heavy metal ions in natural waters have a high mobilization because of their reduction by humic Substances, adsorption, and complexation (Weber, 1988). Organic matter (OM). plays an important role in the accumulation, leaching, and transportation of cations present in water and soil as chelates of extra stability and supplying plant roots with these ions. The equilibrium exchangeable constant has been studied extensively to determine the Stability constant for organic matter-metallic complexes in soils (Kitagishi and Yaman, 1981; Chen et al 2022). Stability constant (SC) values determined by some

Table (1): Solubility constant of metal-HA and FA expressed as logK at different pH value

No.	Cation	pH ₇		pH ₅				pH _{3.5}		pH ₃		
		HA		HA	HA	FA	FA	FA	FA	HA	FA	
1	Cu	12.3	(Kitagishiond and Yaman,1981)									
2	Ni	9.6	12.6	8.7	8.7	4.0	5.8	6.8	6.8	3.3	(Schitzer and Khan,1981)	
3	Co	-	7.6	-	4.1	4.2	3.5	5.4	5.4	3.2	(Kitagishiond and Yaman,1981)	
4	Pb	-	-	-	3.7	4.1	2.2	-	-	2.8	(Schitzer and Khan,1981)	
5	Zn	10.3	-	8.3	6.2	4.0	3.1	-	-	2.7	(Kitagishiond and Yaman,1981)	
6	Mn	5.6	7.2	-	2.3	3.6	1.7	5.1	5.1	2.3	(Schitzer and Khan,1981)	
7	Cd	8.9	0	-	3.8	3.7	1.5	0	0	2.1	(Kitagishiond and Yaman,1981)	
8	Fe	4.8	5.5	6.3	-	-	-	5.3	5.3	-	(Schitzer and Khan,1981)	
9	Ca	6.5	6.4	-	5.8	-	5.1	5.4	5.4	-	(Kitagishiond and Yaman,1981)	
10	Mg	5.5	0	-	2.9	3.4	2.0	0	0	2.7	(Schitzer and Khan,1981)	
11	Fe	6.6	0	-	2.1	2.2	1.2	0	0	1.9	(Kitagishiond and Yaman,1981)	
12	Al	-	8.5	-	-	-	-	11.4	11.4	6.1	(Schitzer and Khan,1981)	
			-	-	-	-	-	-	-	3.7	(Kitagishiond and Yaman,1981)	

authors illustrate the ability of HA and FA to form stable complexes with heavy metal ions (Schnitzer and Kerndorff, 1981). The FA-metal complexes which have lower (SC) are more Soluble and available to plant nutrients, many authors from (3to7) (Table1). Gamble (1986) findings a clearer effect of hydrogen potential (PH) on heavy metal ions sorption on (HA) (Table2). Iron binding (Fe^{+2} and Fe^{+3} ions) in a low pH Solution is very strong and somewhat non-exchangeable with other metal ions. Calcium high stability constant proved that can have a high competition with (Mn) and (Zn) in exchangeable processes (Escrig and Morell, 1998; Yang et al., 2022). Studies on soil organic acid fractionations by Steponova (1974) concluded that there is high affinity between heavy metal ions and (FA). Also Taylor (2003) mentioned that (FA) heavy metal ions content is much higher than found in (HA).

Table (2): Metal Sorption on HA at various pH value percentage of initial metal concentration

No.	Metal	pH _{5,8}	pH _{3,7}	pH _{2,4}
1	Mn	13	3	0
2	Co	45	2	0
3	Zn	64	8	0
4	Cd	77	7	0
5	Cr	100	70	0
6	Ni	61	6	5
7	Al	100	86	7
8	Cu	97	59	12
9	Pb	96	80	19
10	Fe	100	96	81
11	Hg	98	98	99

* Note: After (Gamble, 1986) Initial metal concentration (0.005 mol.L⁻¹)

reported that (SOM) has a low influence on heavy metals uptake by plants (Kördel, 1997). The highest values for (SC) were recorded by Takamatsu and Yoshida. (1978) for Cd^{+2} , Pb^{+2} and Cu^{+2} complexed with humic acid (HA) at pH 5 and that recorded by Kitagishi and Yamane (1981) for Ni^{+2} , Cd^{+2} , Cu^{+2} , and Zn^{+2} at pH7. Andrzejewski and Rosikiewicz (1975) reported that Ni^{+2} , Co^{+2} , and Mn^{+2} complexes with (HA) were soluble, while complexes of Cr^{+3} , Cu^{+2} and Fe^{+2} with humic acid (HA) were insoluble. Augustyn and Urbaniak (1979) observed that Zn^{+2} , Cu^{+2} , and Fe^{+2} are more retained by HA compared to Other heavy metal ions (HM1) such as Al^{+3} and Fe^{+3} which form highly stable complexes with Fulvic acid that interfere easily with Al-hydroxide crystallization (Kodama and Schnitzer, 1980; Yang et al., 2020). Metal complexes Stability with (HA) and (FA) increases as pH increase. Gamble (1986) reported that there are great variable's in the natural system of organic substances in soil and equilibrium function should be determined instead of equilibrium constant. His calculation is based on the average weight of equilibrium function theory, so metal affinity sorption on AH at pH 3.7 take the following order :

$\text{Fe} > \text{Pb} > \text{Cr} > \text{Cu} > \text{Cd} > \text{Zn} > \text{Co}$

Also, Schnitzer and Kerndorff (1981) emphasized the affinity of metal ions to form water-insoluble complexes with FA, this order depends on pH media as follows:

$\text{Fe} = \text{Cr} = \text{Al} > \text{Pb} = \text{Cu} > \text{Hg} > \text{Zn} = \text{Ni} = \text{Cd} = \text{Mn}$

Metal FA complexes are more available to plants compared to those binding with HA which resulted both soluble and insoluble complexes and hydrous oxide. So, HA is relatively in soluble complexes with heavy metals in acid media and considered an organic storage for heavy metals in soils and can regulate heavy metals mobility in soils, (Kabata-penelias, 2000; Abdelrady et al., 2020). FA metal complexes solubility is controlled strongly by FA: metal ratio, so when this ratio is less than (2) water-insoluble complexes formation is favored. According to Bloom and McBride (1979), HA and peat in acid media are likely to bind most divalent cations Zn, Co, Ni, Mn, and Fe as hydrated ions and they conclude that Cu^{+2} ions which in accordance with the functional oxygen group and result from a strong Cu bandit.

Tyler and Olsson (2002) stated that Hg and cd in soft acid media prefer (thiol) legand soft base while acid metals (Mn, Fe) prefer (OH^- , COO^-) hard base legand group and moderate acid metals

such as (Pb, Zn, Cu) tend to form complexes with stronger weak base. In different coal samples Gluskoter, (1977) obtained an index of metal-OM affinities and they distinguished three affinities groups:

Metals of high organic affinities such as B, Br Sb, and Be-

Metals of medium affinity, Ni, Ce, Cr, Cu, and Co -

Metals of low organic affinity but interact with all organic substances, Mn, Mo, Fe, As, and Zn.

Ability of organic matter retention for heavy metals

organic matter (O.M) has a high ability to fix 50% of total heavy metals content in soil and behavior as buffering substances to heavy metal mobility (Trofimova et al. 2017).

Stated that the ability of HA to fix heavy metal cation differs widely between soil horizons and both organic matter acids (HA and FA) to solubilize heavy metals play important role in their cling (Herencia et al, 2008). HA of a soil containing (4%) organic matter may bind about: Fe:17.834, Pb: 4.450 , Zn:1.016 , Mn:0.933 and Cu:1.618(Kg.ha⁻¹) (Halder et al. 2020).

HA ability to complex with heavy metal ions has been Calculated by Vlasov and Michylova (1975) as follows :

Zn: 3:3-Co:3.2 -Cu:3.3 - Mn: 2.6 and Fe: 3.0 gm.Kg⁻¹HA.

According to Rashid (1974), each gm of amino acids in soil may mobilize about 4.5 - 450 mg of different heavy metals with high affinity to Co and Ni and low affinity to Mn. HA sorption Capacity increases as acidity increases, Violante et al (2010) measured the amounts of metals sorbed by HA at pH 5-6 and obtained the following values :

Au: 16- Ce:31-Cd:9-Sr:18 - Cu: 18 and Pb: 120 gm.Kg⁻¹HA.

REFERENCES

- Abdelrady, A., Bachwenkizi, J., Sharma, S., Sefelnasr, A., & Kennedy, M. (2020). The fate of heavy metals during bank filtration: Effect of dissolved organic matter. *Journal of Water Process Engineering*, 38, 101563.
- Al-Hamandi, H. M. N. (2018). Comparison of Chemical Conversion of Humus Compounds in Some Soils of Aqra District. *Tikrit Journal for Agricultural Sciences* , 18(2), 129-134.
- Andrzejewski, M., & Rosikiewicz, D. (1975). Ability of some trace elements to form complexes with humic acids. *Studies about Humus; Transaction of the Internationalsymposium Humus et Planta*.
- Augustyn, D., & Urbaniak, H. (1979). Retention of cations of humic acids from Polish brown coal and some properties of metal-humic compounds. In 7th Int. Symp. Humus and Planta, Brno, Czechoslovakia (p. 26).
- Bloom, P. R., & McBride, M. B. (1979). Metal ion binding and exchange with hydrogen ions in acid-washed peat. *Soil Science Society of America Journal*, 43(4), 687-692.
- Bolan, N. S., & Duraisamy, V. P. (2003). Role of inorganic and organic soil amendments on immobilisation and phytoavailability of heavy metals: a review involving specific case studies. *Soil Research*, 41(3), 533-555.
- Bolton, K. A., & Evans, L. J. (1991). Elemental composition and speciation of some landfill leachates with particular reference to cadmium. *Water, Air, and Soil Pollution*, 60(1), 43-53.
- Boyle, M., & Fuller, W. H. (1987). Effect of municipal solid waste leachate composition on zinc migration through soils (Vol. 16, No. 4, pp. 357-360). American Society of Agronomy, Crop Science Society of America, and Soil Science Society of America.
- Chen, J., Zhang, H., Wei, Q., Farooq, U., Zhang, Q., Lu, T., ... & Qi, Z. (2022). Mobility of water-soluble aerosol organic matters (WSAOMs) and their effects on soil colloid-mediated transport of heavy metal ions in saturated porous media. *Journal of Hazardous Materials*, 129733.
- Escrig, I., & Morell, I. (1998). Effect of calcium on the soil adsorption of cadmium and zinc in some Spanish sandy soils. *Water, Air, and Soil Pollution*, 105(3), 507-520.

- Evans, L. J., Sengdy, B., Lumsdon, D. G., & Stanbury, D. A. (2003). Cadmium adsorption by an organic soil: a comparison of some humic-metal complexation models. *Chemical Speciation & Bioavailability*, 15(4), 93-100.
- Förstner, U., & Müller, G. (2013). *Schwermetalle in Flüssen und Seen als Ausdruck der Umweltverschmutzung*. Springer-Verlag.
- Gamble, D. S. (1986). Interactions between natural organic polymers and metals in soil and freshwater systems: Equilibria. In *The Importance of Chemical "Speciation" in Environmental Processes* (pp. 217-236). Springer, Berlin, Heidelberg.
- Gluskoter, H. J. (1977). Trace elements in coal: occurrence and distribution. Circular no. 499.
- Halder, D., Saha, J. K., & Biswas, A. (2020). Accumulation of essential and non-essential trace elements in rice grain: Possible health impacts on rice consumers in West Bengal, India. *Science of The Total Environment*, 706, 135944.
- Herencia, J. F., Ruiz, J. C., Morillo, E., Melero, S., Villaverde, J., & Maqueda, C. (2008). The effect of organic and mineral fertilization on micronutrient availability in soil. *Soil science*, 173(1), 69-80.
- Kabata-Pendias, A. (2000). *Trace elements in soils and plants*. CRC press. Taylor and Francis group London .
- Killops, V. J., & Killops, S. D. (2013). *Introduction to organic geochemistry*. John Wiley & Sons.
- Kitagishi, K., & Yamane, I. (1981). *Heavy metal pollution in soils of Japan*. Science society press. Tokyo. 302.
- Kodama, H., & Schnitzer, M. (1980). Effect of fulvic acid on the crystallization of aluminum hydroxides. *Geoderma*, 24(3), 195-205.
- Kördel, W. (1997). Fate and effects of contaminants in soils as influenced by natural organic material-status of information. *Chemosphere*, 35(1-2), 405-411.
- Lamy, I., Bourgeois, S., & Bermond, A. (1993). Soil cadmium mobility as a consequence of sewage sludge disposal (Vol. 22, No. 4, pp. 731-737). American Society of Agronomy, Crop Science Society of America, and Soil Science Society of America.
- Laxen, D. P. (1985). Trace metal adsorption/coprecipitation on hydrous ferric oxide under realistic conditions: the role of humic substances. *Water Research*, 19(10), 1229-1236.
- Li, Z., & Shuman, L. M. (1996). Redistribution of forms of zinc, cadmium and nickel in soils treated with EDTA. *Science of the Total Environment*, 191(1-2), 95-107.
- Liu, M., Zhu, J., Yang, X., Fu, Q., Hu, H., & Huang, Q. (2022). Mineralization of organic matter during the immobilization of heavy metals in polluted soil treated with minerals. *Chemosphere*, 301, 134794.
- Matthews, E. (1983). Global vegetation and land use: New high-resolution data bases for climate studies. *Journal of Applied Meteorology and Climatology*, 22(3), 474-487.
- McBride, M. B., Richards, B. K., Steenhuis, T., Russo, J. J., & Sauvé, S. (1997). Mobility and solubility of toxic metals and nutrients in soil fifteen years after sludge application. *Soil Science*, 162(7), 487-500.
- Minkina, T. M., Motuzova, G. V., & Nazarenko, O. G. (2006). Interaction of heavy metals with the organic matter of an ordinary chernozem. *Eurasian Soil Science*, 39(7), 720-726.
- Norvell, W. A., & Lindsay, W. L. (1972). Reactions of DTPA chelates of iron, zinc, copper, and manganese with soils. *Soil Science Society of America Journal*, 36(5), 778-783.
- Pauli, F. W. (1975). Heavy metal humates and their behavior against hydrogen sulfide. *Soil Science*, 119(1), 98-105.
- Rashid, M. A. (1974). Absorption of metals on sedimentary and peat humic acids. *Chemical Geology*, 13(2), 115-123.
- Schnitzer, M., & Kerndorff, H. (1981). Reactions of fulvic acid with metal ions. *Water, Air, and Soil Pollution*, 15(1), 97-108.

- Sholkovitz, E. R., & Copland, D. (1981). The coagulation, solubility and adsorption properties of Fe, Mn, Cu, Ni, Cd, Co and humic acids in a river water. *Geochimica et Cosmochimica Acta*, 45(2), 181-189.
- Stepanova, M. D. (1976). Microelements in the Organic Matter of Chernozems and Soddy-Podzolic Soils.
- Stevenson, F. J. (1972). Organic matter reactions involving micronutrients in soils. *Micronutrients in agriculture*, 79-115.
- Takamatsu, T., & Yoshida, Y. (1978). Determination of stability constants of metal-humic acid complexes by potentiometric titration and ion-selective electrodes. *Soil Science*, 125(6), 377-386.
- Tan, K. H. (2010). *Principles of soil chemistry*. CRC press. Taylor and Francis group London .
- Taylor, T. P., Ding, M., Ehler, D. S., Foreman, T. M., Kaszuba, J. P., & Sauer, N. N. (2003). Beryllium in the environment: a review. *Journal of Environmental Science and Health, Part A*, 38(2), 439-469.
- Trofimova, E. S., Zykova, M. V., Ligacheva, A. A., Sherstoboev, E. Y., Zhdanov, V. V., Belousov, M. V., ... & Dygai, A. M. (2017). Influence of humic acids extracted from peat by different methods on functional activity of macrophages in vitro. *Bulletin of experimental biology and medicine*, 162(6), 741-745.
- Tyler, G., & Olsson, T. (2002). Conditions related to solubility of rare and minor elements in forest soils. *Journal of Plant Nutrition and Soil Science*, 165(5), 594-601.
- Van Dijk, H. (1971). Cation binding of humic acids. *Geoderma*, 5(1), 53-67.
- Violante, A., Cozzolino, V., Perelomov, L., Caporale, A. G., & Pigna, M. (2010). Mobility and bioavailability of heavy metals and metalloids in soil environments. *Journal of soil science and plant nutrition*, 10(3), 268-292.
- Vlasov, N. A., & Mikhailova, A. I. (1975). Interaction of coal humic acid fractions with some metallic cations and the effect of coal humic fertilizers on the distribution of microelements in soil and plants. *Studies about Humus; Transaction of the Internationalsymposium Humus et Planta*.
- Walker, D. J., Clemente, R., Roig, A., & Bernal, M. P. (2003). The effects of soil amendments on heavy metal bioavailability in two contaminated Mediterranean soils. *Environmental Pollution*, 122(2), 303-312.
- Weber, J. H. (1988). Binding and transport of metals by humic materials. *Humic substances and their role in the environment*, 165-178.
- Yang, H. J., Jeong, H. J., Bong, K. M., Kang, T. W., Ryu, H. S., Han, J. H., ... & Na, E. H. (2020). Organic matter and heavy metal in river sediments of southwestern coastal Korea: Spatial distributions, pollution, and ecological risk assessment. *Marine pollution bulletin*, 159, 111466.
- Zunino, H., Aguilera, M., Caiozzi, M., Peirano, P., Borie, F., & Martin, J. P. (1979). Metal-binding organic macromolecules in soil: 3. Competition of Mg (II) and Zn (II) for binding sites in humic and fulvic-type model polymers. *Soil Science*, 128(5), 257-266.

المواد العضوية وامتصاص المعادن الثقيلة

مجبل محمد الجميلي حذيفة معن الحمدي

قسم علوم التربة والموارد المائية / كلية الزراعة / جامعة تكريت/ العراق

الخلاصة

تختلف الترب في محتواها من المادة العضوية والتي تشمل احياء التربة الميتة ونواتج تحللها وبقايا النباتات والمادة الدبالية ثرموديناميكية , تعتبر المادة العضوية غير ثابتة في التربة ولاحقا سوف تتأكسد الى ثاني اوكسد الكربون وماء. من المواد الفعالة الناتجة من تحلل المادة العضوية هما حامض الهيومك وحامض الفولفك واللذان يحتويان على مجاميع فعالة قادرة على اطلاق البروتينات والالكترونيات من خلال تحللها تاركة خلفها جذور كيميائية تتفاعل بصيغة واهبة للإلكترونات اتجاه ايونات العناصر الثقيلة مكونة معقدات دائبة وغير دائبة لكل من حامض الفولفك وحامض الهيومك وهذه المعقدات العضوية المعدنية مختلفة فيما بينها بصفة الثباتية والتي تعتمد بصورة رئيسة على قيمة الاس الهيدروجيني للوسط . اذ ان المعقد الاقل ثباتية هو الاعلى في الذوبانية والحركة في التربة وبالعكس المعقد الاكثر ثباتية هو الاقل ذوبانية وحركة في التربة . لذا تلعب المادة العضوية دورا مهما في التربة من خلال تجمع وغسل ونقل ايونات العناصر الثقيلة الموجودة في المياه والتربة بصورة مخلبية مختلفة الثباتية لتحفيز جذور النباتات بهذه العناصر وتتصرف كأنها مخزن منظم لتجهيز هذه العناصر.

الكلمات المفتاحية:

المخلفات العضوية , المعقدات المعدنية , عامل الثباتية , المعقد المعدني للفولفك اسد , المعقد المعدني للهيومك اسد.