



IRAQI
Academic Scientific Journals



العراقية
المجلات الأكاديمية العلمية

TJAS

Tikrit Journal for
Agricultural
Sciences

ISSN:1813-1646 (Print); 2664-0597 (Online)

Tikrit Journal for Agricultural Sciences

Journal Homepage: <http://www.tjas.org>

E-mail: tjas@tu.edu.iq

Usage of Essential Oils of Natural Herb as Food Preservatives

Yaman Saad Fadhil

Department of Veterinary Public Health, College of Veterinary Medicine, University of Mosul, Mosul, Iraq

* Corresponding author: E-mail: yamansaadds@uomosul.edu.iq

ABSTRACT

One of the most issues that face the food production industries is the manufacture of healthy food products without synthetic preservatives due to the fact that artificial antibacterial agents and chemical additives could have serious detrimental impacts on people's life quality. Essential oils (EOs) extracted from various plant origins have received huge attention because of the possible health benefits. Essential oils are complicated combinations made up of numerous distinct chemicals isolated using various methods. Such chemicals demonstrated substantial biological functions like antioxidant and antibacterial activity through a variety of mechanisms, and they are less dangerous and pose no health problems to humans. In this article, we underline the importance of Essential oils such as major ingredients and sources, antibacterial activities, and potential uses in the food sector. The use of natural additives is becoming more common; nonetheless, they may have negative effects on organoleptic properties. As a result, more research is needed to adjust the dosages used to effectively stop the growth of microbes is still needed.

KEY WORDS:

Essential Oils, Food Preservation, Food Safety, Natural Herbs

Received: 21/07/2022

Accepted: 23/9/2022

Available online: 30/06/2023

© 2023 This is an open access article under the CC by licenses <http://creativecommons.org/licenses/by/4.0>

© 2023 TJAS. College of Agriculture, Tikrit University



استخدام الزيوت الأساسية للنباتات الطبيعية كمواد حافظة للغذاء

يمان سعد فاضل

فرع الصحة العامة البيطرية، كلية الطب البيطري، جامعة الموصل، العراق

الخلاصة

أكثر المشكلات التي تواجه صناعات إنتاج الغذاء هي تصنيع منتجات غذائية صحية بدون مواد حافظة صناعية نظرًا لحقيقة أن العوامل المضادة للبكتيريا الصناعية والإضافات الكيماوية يمكن أن يكون لها آثار ضارة خطيرة على جودة حياة الناس. حظيت الزيوت الأساسية المستخرجة من أصل نباتي باهتمام كبير بسبب الفوائد الصحية المحتملة، حيث أن الزيوت الأساسية تتكون من مكونات معقدة تشمل العديد من المواد الكيميائية المتميزة المعزولة باستخدام طرائق مختلفة، وقد أظهرت هذه المواد الكيميائية المتنوعة ووظائف بيولوجية كبيرة مثل النشاط المضاد للأكسدة والبكتيريا من خلال مجموعة متنوعة من الآليات، وهي أقل خطورة ولا تسبب أي مشاكل صحية للإنسان. تركز هذه المراجعة على أهمية الزيوت الأساسية من حيث

المكونات والمصادر، والأنشطة المضادة للبكتيريا، والاستخدامات المحتملة كمواد حافظة في صناعة الأغذية. أصبح استخدام المضافات الطبيعية أكثر شيوعاً؛ ومع ذلك، قد يكون لها آثار سلبية على الخصائص الحسية. لذلك، لا تزال هناك حاجة إلى مزيد من البحث لضبط الجرعات المستخدمة لوقف نمو الميكروبات بشكل فعال.

الكلمات المفتاحية: الزيوت الأساسية، حفظ الأغذية، سلامة الأغذية، الاعشاب الطبيعية

INTRODUCTION

EOs, commonly referred to as volatile EOs, are aromatic oily substances collected from plants' leaves and peels. Squeezing and steam distillation are two methods for extracting it from plant materials. Steam distillation is widely utilized, especially in commercial production (Lira *et al.*, 2009). The types and concentrations of the substances that make up these oils determine their aroma and flavor, which are different in the oils that are found in many different plants. Additionally, different plants produce varied amounts of EO, which affects the price of the oil. The original dyes help to alter the color of the EO because this can impact how the oils are used in particular foods. EOs are recognized to offer antioxidant and antibacterial effects as food additions. They can also be employed as active ingredients in packing polymers to enhance their characteristics, especially the water vapor barrier feature linked with EOs' hydrophobic nature. Consumer worries about synthetic preservatives have raised interest in organic antibacterial like EOs, which they have food preservation characteristics against a large number of infections (Sonker *et al.*, 2015).

Chemical composition of Essential Oil

Many plants contain EOs, however the part that are the primary source of these oils vary. Plant EOs are typically a mix of polar and non-polar natural ingredients (Masango, 2005). Each oil typically comprises more than a hundred constituents, but the amount of ingredients varies based on the plant that have taken the oil from. However, the most important active chemicals are classified as Terpenoids like (Monoterpenoids and Sesquiterpenoids), and Phenylpropanoids. Both of these categories are formed from distinct main metabolic substrates and are generated via separate metabolic processes. EOs, like all organic chemicals, are formed of hydrocarbon atoms and can be categorized as phenols, terpenes, alcohols, esters, aldehydes, and etc. The two main components of EOs found in a variety of plants and flowers are terpenes and terpenoids. The monoterpenoid and sesquiterpenoid groups of terpenoids provide the most important ingredients of EOs for most plants (Delgoda, 2017).

Almost all EOs contain Monoterpenoids, which have a structure of 10 carbon atoms and at minimum a double bond. Geraniol in lilac, myrcene in hop, linalool in lavender, and pinene in pine are some examples of Monoterpenes (Breitmaier, 2006). Citrus oils are short-lived because they are high in monoterpene hydrocarbons, react quickly with oxygen, and are easily oxidized (Swamy *et al.*, 2015).

Esters are commonly present in a range of EOs and are produced when an alcohol combines with an acid solution (a process named esterification). They have therapeutic properties such as being sedative and antispasmodic. Linalyl acetate is a commonly found ester in lemongrass, lavender and petitgrain EO, also geraniol acetate spotted in sweet marjoram, is among the advantageous components in EOs (Arumugam *et al.*, 2016). Certain esters have antifungal and antibacterial characteristics, such as geranium oil's antifungal effects (Lang and Buchbauer, 2012).

Essential Oils in the food industry

Refrigeration, frozen storage, drying, salting, smoking, and fermentation are all traditional ways of food preservation (Dave and Ghaly, 2011). However, because of the rising desire for low-salt foods, consumers have criticized procedures such as fermentation and salting (Jayasena and Jo, 2013). The meat industry

uses chemical additives like nitrate salt, sulfites, and chlorides to limit the spread of microorganisms that are found in food. These chemicals have a history of causing cancer and other health problems (Jayasena and Jo, 2013). Natural substances have gained popularity as alternatives to artificial preservatives in recent years. In their review Lucera *et al.* (2012) clarified several natural preservatives such as (lactoferrin, lysozyme) derived from animals, bacteriocin from microbial (natamycin, nisin), polymers derived from nature like (chitosan), acidic organic compounds (citric and propionic acid), EOs from plant parts.

Academic researchers have shown that EOs can successfully replace artificial preservatives in various diets (Valková *et al.*, 2021; Ahmed *et al.*, 2021; Radünz *et al.*, 2020). Several studies have looked into the possible benefits of EOs/extracts to increasing the lifespan and inhibiting pathogen development in fresh veggies mixtures (Kraśniewska *et al.*, 2020), fruit juices (Siddiqua *et al.*, 2015), cooked meat (Huq *et al.*, 2015), and turkey (Vasilatos and Savvaidis, 2013). Many papers focused on the usage of EOs in food products, included over 650 studies on fruit, 403 studies on veggies, 415 studies on seafood, 410 studies on beef, 216 studies on milk, and 97 studies on bread and baked products (Fernández-López and Viuda-Martos, 2021). Additional latest studies have looked at the use of rosemary extract in beef (Kaur *et al.*, 2021), the combined effect of EOs in fish conservation (Huang *et al.*, 2021), food packaging (Carpena *et al.*, 2021) and as a natural preservative (Falleh *et al.*, 2020).

Fruits and veggies can be preserved to extend their life span through spraying, soaking, coating, and impregnation (Lucera *et al.*, 2012). A study investigated the impacts of soaking cherry tomatoes in a nanoemulsion of thyme EO against *E. coli*, and the influence of this emulsion combined with ultrasound treatment (He *et al.*, 2021). The investigation revealed that the nanoemulsion by itself can successfully stop the development of *E. coli* on the cherry tomatoes' skin, and the co-treatment had a significant synergistic impact. According to (Kang and Song, 2018), newly cut red mustard leaves that had been rinsed with a nanoemulsion of cinnamon leaf EO had a two-log reduction in the amount of *E. coli*, *L. monocytogenes*, and *S. enterica*. Further investigation from the same researcher revealed that washing with a nanoemulsion of cinnamon leaf EO improves physical separation and inhibits both *Escherichia coli* and *Listeria monocytogenes* on cabbage leaves (Kang *et al.*, 2019). The quality features of mustard (Kang and Song, 2018) and cabbage leaves (Kang *et al.*, 2019) did not demonstrate any negative alterations in the two studies. Lettuce leaves were evaluated after 7 days of storage and found to have reduced *E. coli* levels when washed with a mix of carvacrol - thymol/eugenol comparing to the untreated group (washing with water), as well as unfavorable impacts on sensory qualities (Yuan *et al.*, 2019).

Another study reported the use of a mixture of Spanish oregano oil and Spanish marjoram oil succeeded in inhibiting *listeria monocytogenes* bacteria from a variety of freshly cut veggies without displaying any unfavorable sensory traits (Kraśniewska *et al.*, 2020). According to Dai *et al.* (2021), reported that *Litsea cubeba* EO was applied to cucumbers, carrots, and spinach at the minimum inhibitory doses reduced the number of *Escherichia coli*. The antimicrobial ability of isogenol and coated isogenol against many microbes, fermentation-associated lactic acid bacteria and *Pseudomonas fluorescense* in carrot juice were examined, their investigation found no significant variation in the inhibitory action, contrary to predictions (Nielsen *et al.*, 2017).

Aside from fruits and vegetables, the antibacterial capability of EOs in beef products, particularly meat products, has received considerable attention (Yoo *et al.*, 2021; Khaleque *et al.*, 2016). Pistachio EO (Krichen *et al.*, 2020) and tea tree EO (de Sá Silva *et al.*, 2019) decreased the total number of *L. monocytogenes* in minced meat. The potential of (5, 10%) of clove EO in inhibiting *L. monocytogenes* in minced beef stored at chilling and freezing temperatures were studied, and he noticed that ten percentage of clove EO was a deadly dose for inactivating *L. monocytogenes* regardless of thermal circumstances, but five percentage of clove EO failed to eliminate the pathogen (Khaleque *et al.*, 2016). Neither *E. coli* nor *S. aureus* were significantly reduced in beef as reported by Yoo *et al.* (2021), when used clove EO at (0.5, 1.0 and 1.5%) concentrations. Nonetheless, their research went a step further and revealed that the combination of clove EO with enclosed atmospheric pressure had a bactericidal impact on the two pathogens. Similarly, another study has discovered that chrysanthemum EO mixed with chitosan nanofibers stopped *L. monocytogenes* growth in meat at an

average of almost 100 percent (Lin *et al.*, 2019). A mixture of thyme, cinnamon and clove EO in food matrix was originally used practically by Chaichi *et al.* (2021), the triple mixture at FIC 0.3, 0.39, 0.43 had an impact on *Pseudomonas fluorescein* injected into chicken breast meat, while the mix at a dose of (200 mg/kg) had an immediate antimicrobial impact. As for thyme EO, it effectively stopped the growth of the bacteria *Pseudomonas fluoresce*, *Escherichia coli* and *Salmonella enterica* in ground meat (Jayari *et al.*, 2018). Other research publications indicated that nanoemulsions of EOs efficiently suppressed infections in salmon fillets (Kazemeini *et al.*, 2019) and chicken breast fillets (Noori *et al.*, 2018). In addition to fruits, vegetables, animal products, Eos have been applied and evaluated in baked goods (Khezri *et al.*, 2021; Valková *et al.*, 2021) and dairy products (Ahmed *et al.*, 2021). Table 1 covers some research findings on the application of EOs as a preservative.

Table 1: Latest studies of EOs application as a preserver in the food sector.

Plant source	Food applied	Condition of storage	Findings	References
Garlic EOs	Yogurt	30 days at 4°C	Demonstrated its promise as a natural food preservative against <i>E.coli</i>	Nazari <i>et al.</i> (2019)
<i>Mentha piperita</i>	Stored minced meat	21 days at 4°C	Effective against microbial growth and improved sensory acceptability, prolonging the lifespan of meat by roughly 7 days.	Smaoui <i>et al.</i> (2016)
<i>Rosmarinus officinalis L</i>	Tomato juice	15 days at 5 °C	The antibacterial capabilities of EOs encapsulated by b-CD were preserved during the pasteurization of tomato juice.	Garcia-Sotelo <i>et al.</i> (2019)
Piper betleL	Raw apple juice	15 days at 4°C	In refrigerated circumstances, treated juice increased a 6-day lifetime as compared to untreated juice.	Basak (2018)
<i>Pulicaria inuloides</i>	Fillet Fish	12 days at 10 °C	The results showed that the EOs eradicated the majority of the microorganism. It also extended the lifespan of seafood.	Al-Hajj <i>et al.</i> (2017)

Conclusions and recommendations for future work

In this review, we examined the research on EOs and their significance in the food sector, as well as some of the primary active components, as well as their implementations in the food industry as a preservative technique to prolong the lifespan of foods. It's worth noting that EOs are venturing into a variety of fields. Various studies have shown that several EOs can be utilized as food preservatives, which are often more abundant and have lots of advantages like antibacterial and antioxidant properties, as well as a variety of ways by which they can impact foodborne organisms. The presence of phenolic natural compounds in EOs makes them a vital and healthy option to artificial preservatives and chemical additives, and their presence has proved a substantial ability to resist pathogenic bacteria and other germs without causing nutritional loss.

CONCLUSION

In general, there are numerous obstacles to the applications of EOs in food industry. For instance, antibacterial effect in EOs can be reduced because of many factors that can impact the contents of EOs like geographic location, plant diversity, harvest, way of extraction, and others that can lead to a reduction in EOs implementation. Additionally, it should be noted that EOs have a strong taste and smell, which might alter the flavor and aroma of foods. As a result, investigations should concentrate on the lowest quantity of EO required to retain antibacterial action without altering the organoleptic properties of food products.

CONFLICT OF INTEREST

The authors declare no conflicts of interest associated with this manuscript.

REFERENCES

- Ahmed, L. I., Ibrahim, N., Abdel-Salam, A. B. and Fahim, K. M. (2021). Potential application of ginger, clove and thyme essential oils to improve soft cheese microbial safety and sensory characteristics. *Food Bioscience*, 42, 101177. <https://doi.org/10.1016/j.fbio.2021.101177>.
- Al-Hajj, N. Q. M., Algabr, M., Raza, H., Thabi, R., Ammar, A-F., Aboshora, W. and Wang H. (2017). Antibacterial activities of the essential oils of some aromatic medicinal plants to control pathogenic bacteria and extend the shelf-life of seafood. *Turkish J Fish Aquat Sci.*, 17(1):181–191. https://doi.org/10.4194/1303-2712-v17_1_20.
- Arumugam, G., Swamy, M. K. and Sinniah, U. R. (2016). *Plectranthus amboinicus* (Lour.) Spreng: botanical, phytochemical, pharmacological and nutritional significance. *Molecules*, 21(4), 369. <https://doi.org/10.3390/molecules21040369>.
- Basak, S. (2018). The use of fuzzy logic to determine the concentration of betel leaf essential oil and its potency as a juice preservative. *Food Chemistry*. 240, 1113–1120. <https://doi.org/10.1016/j.foodchem.2017.08.047>.
- Breitmaier, E. (2006). *Terpenes: flavors, fragrances, pharmaca, pheromones*, John Wiley & Sons. <https://doi.org/10.1002/9783527609949>.
- Carpena, M., Nuñez-Estevez, B., Soria-Lopez, A., Garcia-Oliveira, P. and Prieto, M. A. (2021). EOs and their application on active packaging systems: A review. *Resources*, 10(1), 7. <https://doi.org/10.3390/resources10010007>.
- Chaichi, M., Mohammadi, A., Badii, F. and Hashemi, M. (2021). Triple synergistic EOs prevent pathogenic and spoilage bacteria growth in the refrigerated chicken breast meat. *Biocatalysis and Agricultural Biotechnology*, 32, 101926. <https://doi.org/10.1016/j.bcab.2021.101926>.
- Dai, J., Li, C., Cui, H. and Lin, L. (2021). Unraveling the anti-bacterial mechanism of *Litsea cubeba* EO against *E. coli* O157: H7 and its application in vegetable juices. *International Journal of Food Microbiology*, 338, 108989. <https://doi.org/10.1016/j.ijfoodmicro.2020.108989>.
- Dave, D. and Ghaly, A. E. (2011). Meat spoilage mechanisms and preservation techniques: a critical review. *American Journal of Agricultural and Biological Sciences*, 6(4), 486-510. <https://doi.org/10.3844/ajabssp.2011.486.510>.
- De Sá Silva, C., de Figueiredo, H. M., Stamford, T. L. M. and da Silva, L. H. M. (2019). Inhibition of *Listeria monocytogenes* by *Melaleuca alternifolia* (tea tree) EO in ground beef. *International Journal of Food Microbiology*, 293, 79-86. <https://doi.org/10.1016/j.ijfoodmicro.2019.01.004>.

- Delgoda, R. (2017). *Pharmacognosy: Fundamentals, applications and strategies*. Academic Press, 477-494. <https://doi.org/10.1016/b978-0-12-802104-0.00024-x>.
- Falleh, H., Jemaa, M. B., Saada, M. and Ksouri, R. Eos (2020). A promising eco-friendly food preservative. *Food Chemistry*, 330, 127268. <https://doi.org/10.1016/j.foodchem.2020.127268>.
- Fernández-López, J. and Viuda-Martos, M. (2018). Introduction to the special issue: application of EOs in food systems. *Foods*, 7(4), 56. <https://doi.org/10.3390/foods7040056>.
- García-Sotelo, D., Silva-Espinoza, B., Perez-Tello, M., Olivas, I., Alvarez-Parrilla, E., González-Aguilar, G. A. and Ayala-Zavala, J. F. (2019). Antimicrobial activity and thermal stability of rosemary essential oil: β -cyclodextrin capsules applied in tomato juice. *LWT*, 111:837–845. <https://doi.org/10.1016/j.lwt.2019.05.061>.
- He, Q., Guo, M., Jin, T. Z., Arabi, S. A. and Liu, D. (2021). Ultrasound improves the decontamination effect of thyme EO nanoemulsions against *Escherichia coli* O157: H7 on cherry tomatoes. *International Journal of Food Microbiology*, 337, 108936. <https://doi.org/10.1016/j.ijfoodmicro.2020.108936>.
- Huang, X., Lao, Y., Pan, Y., Chen, Y., Zhao, H., Gong, L., ... and Mo, C. H. (2021). Synergistic antimicrobial effectiveness of plant EO and its application in seafood preservation: A review. *Molecules*, 26(2), 307. <https://doi.org/10.3390/molecules26020307>.
- Huq, T., Vu, K. D., Riedl, B., Bouchard, J. and Lacroix, M. (2015). Synergistic effect of gamma (γ)-irradiation and microencapsulated antimicrobials against *Listeria monocytogenes* on ready-to-eat (RTE) meat. *Food microbiology*, 46, 507-514. <https://doi.org/10.1016/j.fm.2014.09.013>.
- Jayari, A., El Abed, N., Jouini, A., Mohammed Saed Abdul-Wahab, O., Maaroufi, A. and Ben Had j., Ahmed S. (2018). Antibacterial activity of *Thymus capitatus* and *Thymus algeriensis* EOs against four food-borne pathogens inoculated in minced beef meat. *Journal of Food Safety*, 38(1), e12409. <https://doi.org/10.1111/jfs.12409>.
- Jayasena, D. D. and Jo, C. (2013). Essential oils as potential antimicrobial agents in meat and meat products: A review. *Trends in Food Science & Technology*, 34(2), 96-108. <https://doi.org/10.1016/j.tifs.2013.09.002>.
- Kang, J. H. and Song, K. B. (2018). Inhibitory effect of plant EO nanoemulsions against *Listeria monocytogenes*, *Escherichia coli* O157: H7, and *Salmonella Typhimurium* on red mustard leaves. *Innovative Food Science and Emerging Technologies*, 45, 447-454. <https://doi.org/10.1016/j.ifset.2017.09.019>.
- Kang, J. H., Park, S. J., Park, J. B. and Song, K. B. (2019). Surfactant type affects the washing effect of cinnamon leaf EO emulsion on kale leaves. *Food chemistry*, 271, 122-128. <https://doi.org/10.1016/j.foodchem.2018.07.203>.
- Kaur, R., Gupta, T. B., Bronlund, J. and Kaur, L. (2021). The potential of rosemary as a functional ingredient for meat products-a review. *Food Reviews International*, 1-21. <https://doi.org/10.1080/87559129.2021.1950173>.
- Kazemeini, H., Azizian, A. and Shahavi, M. H. (2019). Effect of chitosan nano-gel/emulsion containing *bunium persicum* EO and nisin as an edible biodegradable coating on *escherichia coli* o157: H7 in rainbow trout fillet. *Journal of Water and Environmental Nanotechnology*, 4(4), 343-349.
- Khaleque, M. A., Keya, C. A., Hasan, K. N., Hoque, M. M., Inatsu, Y. and Bari, M. L. (2016). Use of cloves and cinnamon EO to inactivate *Listeria monocytogenes* in ground beef at freezing and refrigeration temperatures. *LWT*, 74, 219-223. <https://doi.org/10.1016/j.lwt.2016.07.042>.

- Khezri, S., Khezerlou, A. and Dehghan, P. (2021). Antibacterial activity of *Artemisia persica* Boiss EO against *Escherichia coli* O157: H7 and *Listeria monocytogenes* in probiotic Doogh. *Journal of Food Processing and Preservation*, 45(5), e15446. <https://doi.org/10.1111/jfpp.15446>.
- Kraśniewska, K., Kosakowska, O., Pobiega, K. and Gniewosz, M. (2020). The influence of two-component mixtures from Spanish *Origanum* oil with Spanish Marjoram oil or coriander oil on antilisterial activity and sensory quality of a fresh cut vegetable mixture. *Foods*, 9(12), 1740. <https://doi.org/10.3390/foods9121740>.
- Krichen, F., Hamed, M., Karoud, W., Bougatef, H., Sila, A. and Bougatef, A. (2020). EO from pistachio by-product: Potential biological properties and natural preservative effect in ground beef meat storage. *Journal of Food Measurement and Characterization*, 14(6), 3020-3030. <https://doi.org/10.1007/s11694-020-00546-6>.
- Lang, G. and Buchbauer, G. (2012). A review on recent research results (2008–2010) on essential oils as antimicrobials and antifungals. A review. *Flavour and Fragrance Journal*, 27(1), 13-39. <https://doi.org/10.1002/ffj.2082>.
- Lin, L., Mao, X., Sun, Y., Rajivgandhi, G. and Cui, H. (2019). Antibacterial properties of nanofibers containing chrysanthemum EO and their application as beef packaging. *International journal of food microbiology*, 292, 21-30. <https://doi.org/10.1016/j.ijfoodmicro.2018.12.007>.
- Lira, P. D. L., Retta, D., Tkacik, E., Ringuelet, J., Coussio, J. D., Van Baren, C. and Bandoni, A. L. (2009). Essential oil and by-products of distillation of bay leaves (*Laurus nobilis* L.) from Argentina. *Industrial Crops and Products*, 30(2), 259-264. <https://doi.org/10.1016/j.indcrop.2009.04.005>.
- Lucera, A., Costa, C., Conte, A. and Del Nobile, M. A. (2012). Food applications of natural antimicrobial compounds. *Frontiers in microbiology*, 3, 287.
- Masango, P. (2005). Cleaner production of EOs by steam distillation. *Journal of Cleaner Production*, 13(8), 833-839. <https://doi.org/10.1016/j.jclepro.2004.02.039>.
- Nazari, M., Ghanbarzadeh B., Kafil H. S., Zeinali M. and Hamishehkar H. (2019). Garlic essential oil nanophytosomes as a natural food preservative: its application in yogurt as food model. *Colloid Interface Science Communications*, 30, 100176. <https://doi.org/10.1016/j.colcom.2019.100176>.
- Nielsen, C. K., Kjems, J., Mygind, T., Snabe, T., Schwarz, K., Serfert, Y. and Meyer, R. L. (2017). Antimicrobial effect of emulsion-encapsulated isoeugenol against biofilms of food pathogens and spoilage bacteria. *International journal of food microbiology*, 242, 7-12. <https://doi.org/10.1016/j.ijfoodmicro.2016.11.002>.
- Noori, S., Zeynali, F. and Almasi, H. (2018). Antimicrobial and antioxidant efficiency of nanoemulsion-based edible coating containing ginger (*Zingiber officinale*) EO and its effect on safety and quality attributes of chicken breast fillets. *Food control*, 84, 312-320. <https://doi.org/10.1016/j.foodcont.2017.08.015>.
- Radünz, M., dos Santos Hackbart, H. C., Camargo, T. M., Nunes, C. F. P., de Barros, F. A. P., Dal Magro, J., ... and da Rosa Zavareze, E. (2020). Antimicrobial potential of spray drying encapsulated thyme (*Thymus vulgaris*) essential oil on the conservation of hamburger-like meat products. *International Journal of Food Microbiology*, 330, 108696. <https://doi.org/10.1016/j.ijfoodmicro.2020.108696>.
- Siddiqua, S., Anusha, B. A., Ashwini, L. S. and Negi, P. S. (2015). Antibacterial activity of cinnamaldehyde and clove oil: effect on selected foodborne pathogens in model food systems

- and watermelon juice. *Journal of food science and technology*, 52(9), 5834-5841. <https://doi.org/10.1007/s13197-014-1642-x>.
- Smaoui S., Hsouna A. B., Lahmar A., Ennouri K., Mtibaa-Chakchouk A., Sellem I., Najah S., Bouaziz M. and Mellouli L. (2016). Bio-preservative effect of the essential oil of the endemic *Mentha piperita* used alone and in combination with BacTN635 in stored minced beef meat. *Meat Science*, 117,196–204. <https://doi.org/10.1016/j.meatsci.2016.03.006>.
- Sonker, N., Pandey, A. K. and Singh, P. (2015). Efficiency of *Artemisia nilagirica* (Clarke) Pamp. essential oil as a mycotoxicant against postharvest mycobiota of table grapes. *Journal of the Science of Food and Agriculture*, 95(9), 1932-1939. <https://doi.org/10.1002/jsfa.6901>.
- Swamy, M. K., Mohanty, S. K., Sinniah, U. R. and Maniyam, A. (2015). Evaluation of patchouli (*Pogostemon cablin* Benth.) cultivars for growth, yield and quality parameters. *Journal of essential oil bearing plants*, 18(4), 826-832. <https://doi.org/10.1080/0972060x.2015.1029989>.
- Valková, V., Ďúranová, H., Galovičová, L., Vukovic, N. L., Vukic, M. and Kačániová, M. (2021). In Vitro antimicrobial activity of lavender, mint, and rosemary essential oils and the effect of their vapours on growth of *Penicillium* spp. in a bread model system. *Molecules*, 26(13), 3859. <https://doi.org/10.3390/molecules26133859>.
- Vasilatos, G. C. and Savvaidis, I. N. (2013). Chitosan or rosemary oil treatments, singly or combined to increase turkey meat shelf-life. *International journal of food microbiology*, 166(1), 54-58. <https://doi.org/10.1016/j.ijfoodmicro.2013.06.018>.
- Yoo, J. H., Baek, K. H., Heo, Y. S., Yong, H. I. and Jo, C. (2021). Synergistic bactericidal effect of clove oil and encapsulated atmospheric pressure plasma against *Escherichia coli* O157: H7 and *Staphylococcus aureus* and its mechanism of action. *Food Microbiology*, 93, 103-611. <https://doi.org/10.1016/j.fm.2020.103611>.
- Yuan, W., Teo, C. H. M. and Yuk, H. G. (2019). Combined antibacterial activities of EO compounds against *Escherichia coli* O157: H7 and their application potential on fresh-cut lettuce. *Food Control*, 96, 112-118.