



IRAQI
Academic Scientific Journals



العراقية
المجلات الأكاديمية العلمية

TJAS

Tikrit Journal for
Agricultural
Sciences

ISSN:1813-1646 (Print); 2664-0597 (Online)

Tikrit Journal for Agricultural Sciences

Journal Homepage: <http://www.tjas.org>

E-mail: tjas@tu.edu.iq

Improving Vitamins C and D₃ Content of Medicinal Pumpkin Plant Grown under Organic Farming Programs

Taha S. Ahmed¹, Bnar J. Jalal², Rajaa M. Hasaballa³, and Mustafa R. Alqaisi⁴

^{1,4} College of Agriculture - Tikrit University, Iraq

² College of Agricultural Engineering Science - University of Sulaimani, Iraq

³ College of Education For women Tikrit University, Iraq

*Corresponding author: E-mail: taha.shaheb@tu.edu.iq

ABSTRACT

KEY WORDS:

organic fertilizers, organic farming, farming programs, spent mushroom substrate, vitamins C and D₃.

Received: 30/01/2024
Revision : 29/04/2024
Proofreading: 06/05/2024
Accepted: 23/05/2024
Available online: 30/06/2024

© 2024. This is an open access article under the CC by licenses <http://creativecommons.org/licenses/by/4.0>



This study aimed to evaluate the application of different farming programs. Three organic farming (P₂, P₃, and P₄) were used and compared with traditional program P₁ (chemical). Their effect in improving medicinal pumpkin seeds quality was examined. As a quality addition of supplying chain, the quantitative and qualitative content of vitamins C and D₃ were assayed. This plant seeds are considered of healthy and medical importance, and have their marketing placement. A big amount of the seeds has been consumed directly by human year round. Using the seeds in food production processes increased their nutritional value. The field experiment was conducted on two dates. The design used was the randomized complete block design. All data were collected and analyzed according to ANOVA table using Duncan's multiple range test at P= 0.05. The results obtained showed significant distinguish of the organic farming program (P₃) compared to the rest the programs used. This treatment had significant increase of vitamins C and D₃ content in the seeds in the two dates (0.0883 mg l⁻¹, 0.08671 mg l⁻¹, 12.38 mg l⁻¹ and 12.32 mg l⁻¹ respectively). The other two organic programs (P₂, P₄) were significantly distinct compared to the traditional one (P₁). The study concluded that these studied farming programs can be an effective choice to improve seed quality of this crop. The healthy and nutritional value are enhanced, and the side effects of traditional agriculture on the environment are reduced.

تحسين محتوى فيتاميني C وD₃ في نبات القرع الطبي المزروع تحت برامج الزراعية العضوية

طه شهاب احمد¹، بنار جمال جلال²، رجاء منصور حسب الله³ ومصطفى رشيد القيسي⁴

⁴1 كلية الزراعة-جامعة تكريت

² كلية علوم الهندسة الزراعية-جامعة السليمانية

³ كلية التربية للبنات-جامعة تكريت

الخلاصة

هدفت هذه الدراسة إلى تقييم مدى تطبيق برامج زراعية مختلفة. تم استخدام ثلاثة برامج للزراعة العضوية (P₂ و P₃ و P₄) ومقارنتها ببرنامج للزراعة التقليدية (P₁ الكيمائي). تم دراسة تأثيرها في تحسين جودة بذور القرع الطبي. وكإضافة نوعية لسلسلة التوريد الغذائي، تم تقدير المحتويات الكمية والنوعية لفيتاميني C وD₃، تعد بذور هذا النبات ذات أهمية صحية وطبية، ولها مكانتها التسويقية. يتم استهلاك كمية كبيرة من البذور مباشرة من قبل الإنسان على مدار السنة. وقد أدى استخدام البذور في عمليات الانتاج الغذائي إلى زيادة قيمتها الغذائية. أجريت التجربة الحقلية في مواعدين اثنتين. وكان التصميم المستخدم هو تصميم القطاعات العشوائية الكاملة. جمعت البيانات وحللت وفقاً لجدول ANOVA باستخدام اختبار Duncan متعدد المدى عند مستوى احتمالية P = 0.05. أظهرت النتائج تميزاً معنوياً لبرنامج الزراعة العضوية (P₃) مقارنة ببقية البرامج المستخدمة. أدت هذه المعاملة إلى زيادة معنوية في محتوى فيتاميني C وD₃ في البذور عند المواعدين المستخدمين (0.0883 ملغم لتر⁻¹، و0.08671 ملغم لتر⁻¹ وكذلك 12.38 ملغم لتر⁻¹ و12.32 ملغم لتر⁻¹ على التوالي). كان البرنامجين العضويين الآخرين (P₂ و P₄) مختلفين بشكل كبير مقارنة بالبرنامج التقليدي (P₁). وخلصت الدراسة إلى أن هذه البرامج العضوية المدروسة يمكن أن تكون خياراً فعالاً لتحسين نوعية بذور هذا المحصول. حيث تبرز القيمة الصحية والغذائية، وتقلل الآثار الجانبية للزراعة التقليدية على البيئة.

الكلمات المفتاحية: الأسمدة العضوية، الزراعة العضوية، البرامج الزراعية، ركيزة الفطر المستهلك، فيتامينات C وD₃.

INTRODUCTION

Styrian pumpkin variety was first appeared in Southeastern Austria Styria (Fruhvirth and Hermetter, 2008). Common pumpkin (*Cucurbita pepo* convar. *pepo* var. *styriaca*) is annual plant related to the family of Cucurbitaceae. It has extended sprawling prickly stem, nearly 300 - 500 cm, with female flowers on stalks and male flowers appeared in leaf axil. Also, the plant has fibrous root and fruits colored green to orange (Aroiee and Omidbaigi, 2002). Styria is an economically significant horticultural plant. It is widely grown for its valuable seeds that contain protein and minerals. It is well known that oil extracted from the seeds is low in cholesterol. This material, in men, has an effect in reducing symptoms of benign prostatic hyperplasia. In women, it is decreasing hot flashes, reliving hormone-related headaches, and easing blood pressure. Fighting heart diseases, treating overactive bladder, and reducing hair loss are other benefits (Ebrahimzadeh, et al., 2021 and Al-Jaf et al., 2023).

Seed of pumpkin is rich source of oils essential for human body. Namely, they contain several nutraceuticals such as β -sitosterol, amino acids, unsaturated fatty acids, tocopherols, phenolic compounds, and cucurbitacins as active substances (Kim et al., 2012). Therefore, they are used in manufacturing various drugs such as Grunfig, Pepostrin, and Peponen. Medications for gastrointestinal regulation, atherosclerosis, and urinary tract inflammation are also included (Fadaei and Akbarpour, 2023). In addition, they are rich in omega-3, 6 and 9. Seed oil can promote wellness of human immunodeficiency virus\ acquired immunodeficiency syndrome, HIV\ AIDS (Najafi et al., 2021). Moreover, it has antioxidant

properties due to presence of some vitamins such as C and β -carotene included in vitamin A (Jeznach *et al.*, 2012).

Recently, vitamin D₃ deficiency has been one of the important issues people face. The vitamin is mostly derived from animal sources. Thus, using nutritional supplements and chemical medications by vegetarian people is increased, especially during the COVID-19 period. Researchers have attempted to discover different sources other than animal or chemical to obtain this vitamin. Some research reported that D₃ can be available in some plants belonging to Solanaceae and Cucurbitaceae families (Aburjai, *et al.*, 1998, Japelt, 2011, and Black, *et al.*, 2017; Al-Fhdawi *et al.*, 2023). Vitamin C is an effective safe nutrients. Vitamin C an importance role in human health such as protection against immune system deficiencies, prenatal health problems, cardiovascular disease. Pumpkin seed content of vitamin C 15.76-26.02 mg 100g⁻¹ (Barros *et al.*, 2024).

Lately, several studies have been conducted on secondary metabolites of medicinal plants to increase their natural components and yield. In the farming of medicinal plants management, applying materials of minimal damaging side effects on human health and environment is highly recommended (Fadaei and Akbarpour, 2023). This can be achieved under organic farming programs depending on the use of organic and bio fertilizers, ash, bio-controlling, and plant extraction. This technique protects the environment and supports sustainability. Furthermore, Raw materials of such plants are of high quality and free from harmful chemical residues including pesticides, fertilizers, and heavy metals (Ahmed, *et al.*, 2022).

Organic fertilizers improve soil texture and regulate pH value. Eventually, yield components, active substances, and mineral composition of the plant are influenced. Nitrogen (N) is one of the components of poultry manure. This element improves soil productivity by increasing organic carbon and microorganisms (Hialy, 2018). Poultry manure also contains nutrients essential for plant's growth, such as calcium, magnesium, sulphur, manganese, copper, zinc, iron, boron and molybdenum (Suhartini, *et al.*, 2020). Spent mushroom substrate (SMS) is one of these fertilizers used in plant production. It has essential role through amending soil properties; especially supplying minerals and organic materials. Consequently, leads to an increase in plant growth, quantitative and qualitative yield characteristics (Ahmed, *et al.*, 2023).

Compost tea has a great importance in sustainable agriculture as well. It directly provides mineral nutrients to the plant (Ros, *et al.*, 2020). Bio fertilizers comprise variety of useful microorganisms such as *Azotobacter* and *Trichoderma*. This type improves and accelerates plant growth and protects plants from diseases (Habibi, *et al.*, 2015). The high quantity and quality medicinal plants can be produced under ideal growth requirements, particularly environmental conditions. Planting date is one crucial factor; as early sowing during spring may expose seedlings to cold and possible frost damage. On the other hand, delaying planting may negatively affect plant growth and development. Exposure to unsuitable environmental conditions causes lack growth and less accumulation of active compounds in plants (Omidbeigi, 2004 and Latifi, *et al.*, 2012). To achieve best value of active compound such as antioxidant, phenolic substances, and flavonoids in pumpkin, planting time is strongly considered (Oloyede, *et al.* 2014 and Ahmed, *et al.* 2022). According to the importance of Styrian pumpkin and farming programs, this study was proposed. The aim of the study was to evaluate the efficiency of organic farming programs on

vitamins C and D₃ in this plant. In addition, investigating the effect of sowing date was examined.

MATERIALS AND METHODS

This experiment was conducted in the research center at the Department of Horticulture and landscape, Tikrit University. Medicinal pumpkin plants (*Cucurbita pepo* Subsp. *pepo* Var. *Styriaca*) was examined in this experiment. Analysis of experiment soil was implemented (table 1). Four treatments (terraces) were applied with three replications. On both sides of each terrace, ten plants were planted for each treatment. Each experimental unit was 2.5*0.7 m dimensions. This study was one-way experiment. The treatments included four farming programs; chemical farming program as a control P₁, organic P₂, organic P₃, and organic P₄. Each of which was implemented on March 10,2019 and April 1, 2019 (table 2). Properties of the organic fertilizers used in the programs 2,3, and 4 are shown in table 3.

Table 1. Physical and chemical properties of field soil

Measurements	Value	Measurements	Value %
N	6 ppm	Gypsum	16.3
K	40.05 ppm	Lime	21.4
Na	134.5 ppm	O.M	1.2
P	9.5 ppm	Sand	58
EC	4.475 ms cm ⁻¹	Loam	24
pH	8	Clay	18
Texture	sandy loam		

Table 2. Weather data for the experiment location (2019)

Date Month	Solar Mj/m2/m	Min. Humidity %	Max. Humidity %	Daily low Temp. °C	Daily high Temp. °C	Temp. average °C	Min. Temp. °C	Max. Temp. °C	Rain mm
March	15.43	40.21	91.70	2.39	24.86	14.04	7.57	20.50	46.40
April	18.27	23.49	75.68	7.74	30.20	17.81	11.36	24.25	33.00
May	24.61	18.20	57.72	11.57	45.92	28.43	19.45	37.41	6.40
Jun	27.28	10.51	28.63	22.26	48.83	35.47	26.86	44.09	0.00
July	27.94	15.96	30.58	22.68	46.33	35.42	26.80	44.03	0.00

Table 3. The properties of organic fertilizers

Measurement	Unit	Organic fertilizer		Measurement	Unit	Organic fertilizer	
		SMS	poultry manure			SMS	poultry manure
N	ppm	9016	17710	EC	mS.cm ⁻¹	5.2	4.85
K	ppm	1550	3200	pH	—	8.1	7.8
Na	ppm	7350	2700	O.M	%	24.6	19.2
P	ppm	23.5	11	O.C	%	14.3	11.2
C/N ratio		1/16	1/6				

The studied programs:

1- Program 1 (P₁) included ground addition of compound (NPK 12-8-16 + TE) and foliar addition of compound 20-20-20 + TE + Amino acid. Chemical controlling with Malathion - 50 wp (g L⁻¹) was used. This treatment was applied as a control.

2- Program 2 (P₂) represented applying 16 t h⁻¹ spent mushroom substrate (SMS) for all plants in this experimental unit. 1.5 g ash, 1 g *Trichoderma sp.* (19*10⁷ spore g⁻¹), *Azotobacter sp.* 10 ml (2*10⁹ CFU ml⁻¹) were separately added to each planting hole of this treatment. In addition, foliar spraying of compost tea processed from SMS was added on the plant shoots. Six weeks after first sowing date, 10 ml plant⁻¹ of *Trichoderma* was added. After four weeks, the process of bio-controlling was repeated. Then, extractions of garlic and eucalyptus were applied to the plants.

3- Program 3 (P₃) comprised of the same applications involved in program 2 with moderation. Ground and foliar additions of SMS excluded. Instead, poultry manure was added.

4- In the program P₄ (1:1 SMS and poultry manure), 50% of each P₃ and P₄ were applied. All farm practices and application dates are displayed in table 4.

Table 4. Application dates of agricultural practices used in the experiment.

Agricultural practices	Applying Dates	
	Date one	Date two
Sowing date	10/3/2019	1/4/2019
Ground applying fertilizers	23/2/2019	18/3/2019
Biofertilizers	Trichoderma	24/2/2019
		19/3/2019
		24/3/2019
Foliar application	Azotobacter	19/4/2019
	First	3/3/2019
	Second	25/3/2019
Controlling applications	Third	3/4/2019
	Trichoderma	9/4/2019
	Plants	25/4/2019
Garlic extraction	Garlic	15/5/2019
	eucalyptus	10/5/2019
	Chemical	10/5/2019
Ash	24/4/2019	5/5/2019
	24/5/2019	30/5/2019
	20/4/2019	10/5/2019
	20/4/2019	10/5/2019
	20/4/2019	10/5/2019
	23/2/2019	18/3/2019

The seed content of ascorbic acid (vitamin C) and vitamin D₃ were measured in this experiment (Kozhanova, et al., 2002). HPLC model SYKAM (Germany) provided with vitamins detector UV- 280 nm. The mobile phase was an isocratic acetonitrile: D.W: (75: 25)

at flow rate at 0.7 mL/min, column was C18 – ODS (25 cm * 4.6 mm). Randomized Complete Block Design (RCBD) was used for this one-way experiment. All results were statistically analyzed using GenStat 12th ed. According to Payne, *et al.* (2009). Duncan's multiple range test was implemented to compare the mean values at $P= 0.05$.

RESULTS AND DISCUSSION

Farming programs impacted vitamin C content positively in the date one (figure 1) and the date two (figure2). The result showed that program P₃ significantly higher in vitamin C content compared with P₄ (25%), P₂ (50%), and P₁ (59%). Farming Program P₁ was significantly lowest compared with the other farming programs (figure 1).




Figure 1. Effect of farming programs on vitamin C in medicinal pumpkin plant under date 1

Also, the farming program P₃ responded similarly in the date two. Significantly the highest vitamin C was observed in P₃. Vitamin C reduced in P₄ (37%), P₂ (53%), and P₁ (50%) compared with P₃ (figure 2). According to statically analysis, there were no significant different between P₂ (0.04355 mg l⁻¹) and P₁ (0.03078 mg l⁻¹). Vitamin C content was inconstant between the date one and the date two.




Figure 2. Effect of farming programs on vitamin C in medicinal pumpkin plant under date 2

Figure (3) explained the effect of farming programs on vitamin D₃ on date 1. The result showed that program P₃ caused significant effect on the D₃. This treatment gave the highest value (12.38 mg l⁻¹), compared to the lowest (7.77 mg l⁻¹) under the treatment of program P₂.

The different in planting date with different environment condition (table 2) might be play an important role for changing the amount of vitamin D₃.




Figure 3. Effect of farming programs on vitamin D₃ in medicinal pumpkin plant under date 1

The results in figure (4) exhibited the effect of farming programs on vitamin D₃ under date 2. The program P₃ significantly influenced the concentration of vitamin D₃. This parameter reached the highest record (12.32 mg l⁻¹) in program P₃. In comparison, the

parameter reached the lowest value (6.71 mg l⁻¹) in the which scored when compared to program P₂.




Figure 4. Effect of farming programs on vitamin D₃ in medicinal pumpkin plant under date 2

The result showed that program P₃ had significant increment on the concentration of vitamin C on the dates 1 and 2. The reason of this increase may be due to the role played by poultry manure used in this treatment. This fertilizer is rich in macro-nutrients such as nitrogen, phosphor and potassium (table 3). Nitrogen promotes plant development and improves physiological characteristics, which affects the growth, yield, and quality of plant parts (Liu *et al.*, 2023). Phosphorus plays a crucial role in the formation of cellular membranes and regulates metabolic processes, and promotes plant growth and physiological metabolism (Afzal *et al.*, 2023). Potassium activates various enzymes involved in intracellular osmotic regulation, membrane protein transport, and photosynthesis transport. These are fundamental for plant growth and development (Khandaker, *et al.*, 2022; Fadam and Hamad, 2023)

On other hand, it was confirmed that this small quantities of vitamin C, presented relatively in pumpkin seeds, are subject to plant genotype and growing conditions (Jeznach, *et al.*, 2012). This result was agreed with Isuwan (2014). Concentration of vitamin C in date 1 was increased compared to date 2. Another reason for this increment may be because of compost tea which contains mineral nutrients provided in available forms for the plant. Because it is a ready and instant resource for mineral nutrition, the compost in this program (P₃), had a direct effect in plant nutrition. This caused positive effects on plant growth and productivity. Ultimately, vitamin C was positively affected (Hirzel, *et al.*, 2012).

Increase vitamin D₃ in program (P₃) under the dates 1 and 2 (figures 3, 4) may be due to poultry manure (Baur, *et al.*, 2016). This material had effect in improving soil structure and ventilation. The fertilizer is also important for soil microorganisms. In addition, - have roles in the forming of organic acids and plant growth. The good content of N, P, K, and C\N ration in poultry manure (table 3), give the preference of P₃ to increase the vitamins in plant seeds. This may enhanced C and D₃ content (Ahmed, *et al.*, 2022).

All material included in program (P₃) had an effect to rise metabolism in plant which increased concentrations of vitamins C and D₃. The reason of accumulating these vitamins could be because lifespan of the plants in date 1 was longer than that of date 2. Accordingly, C and D₃ increased in the plants exposed to higher light intensities (Oyetunde, *et al.*, 2023). Or maybe the environmental conditions were more suitable in date 1 during plant growth stages and development (table 2).

CONCLUSION

Styrian Pumpkin, *Cucurbita pepo* ssp. *pepo* var. *styriaca* is considered one of the important plants. Organic program P₃ was superior in most characteristics studied. that showed the distinctive role of organic farming programs in sustaining the health of the environment and biological diversity. Even though, there was a short period between the planting dates, there were some variation in the vitamin C and D. Accordingly, the use of organic farming program is recommended. In addition, use of other agricultural practices that contribute to improve approaches environmentally safe can be adopted.

REFERENCES

- Aburjai, T, Suleiman A and Mustafa A. (1998). Vitamin D₃ and its metabolites in tomato, potato, eggplant and zucchini leaves. *Phytochemistry*. Elsevier Science Ltd. All rights reserved Printed in Great Britain. Vol. 49, No. 8, pp. 2497±2499.
- Afzal, A., Bahader, S., Ul Hassan, T., Naz, I., & Din, A. U. (2023). Rock phosphate solubilization by plant growth-promoting *Bacillus velezensis* and its impact on wheat growth and yield. *Geomicrobiology Journal*, 40(2), 131-142.
- Ahmed, T.S., A.H. Saeed, G.J. Zaidan, and M.R.M. Alqaisi (2023). Effect of spent mushroom substrate and sulfur on growth and yield traits of broccoli in gypsiferous soil. *Sabrao Journal of Breeding and Genetics*. 55 (2) 564-574.
- Ahmed, T.S., Alqaisi, M.R. and Zahwan, T.A. (2022). Evaluation of seed yield , oil , and fatty acid ratio in medicinal pumpkin grown according to traditional cultivation and different programs of organic farming, *Organic Agriculture* [Preprint], (0123456789). doi.org/10.1007/s13165-021-00377-9.
- Aroiee, H. and Omidbaigi, R. (2002). Effects of pre- and post-emergence salt treatment on some characteristics of medicinal pumpkin (*Cucurbita pepo* convar. *pepo* var. *styriaca*)' *Acta Horticulturae*, 576, pp. 253–257. doi.org/10.17660/ActaHortic.2002.576.36.
- Al-Jaf, H. M. ; Mubarak, I., K. ; Abed, A., M. ; & Al-Dulaimy, F. Z. A. (2023). Artificial seeds technology: a review. *Anbar journal of Agricultural Sciences*, 21(2), 396-407.

doi: 10.32649/ajas.2023.181869

- Al-Fhdawi, E. H. O., Al-Ubaidi, O. M., & Mahmood, N. J. (2023). Genotypic and phenotypic variances and correlations as affected by planting and popping in Zea mays Avarta. *Anbar Journal of Agricultural Sciences*, 21(2), 599-612. doi: 10.32649/ajas.2023.179765
- Barros, S. L., Frota, M. M., de Menezes, F. L., de Brito Araújo, A. J., dos Santos Lima, M., Fernandes, V. B., ... & de Vasconcelos, L. B. (2024). Physical–chemical, functional and antioxidant properties of dehydrated pumpkin seeds: Effects of ultrasound time and amplitude and drying temperature. *Waste and Biomass Valorization*, 15(2), 1123-1140.
- Baur, A.C., Brandsch, C., König, B., Hirche, F. and Stangl, G.I. (2016). Plant Oils as Potential Sources of Vitamin D, *Frontiers in Nutrition*, 3(August). doi.org/10.3389/fnut.2016.00029.
- Black, Lucinda J., Robyn, M. Lucas, Jill, L. Sherriff, Lars, Olof Björn and Janet, F. Bornman (2017). In Pursuit of vitamin d in plants. *nutrients*, 9, 136; doi:10.3390/nu9020136.
- Ebrahimzadeh, H., Lotfi, M. and Sadat-Hosseini, M. (2021). Parthenogenetic Haploid Plant Production in Styrian Pumpkin by Gamma Irradiated Pollen, *International Journal of Horticultural Science and Technology*, 9(2), pp. 305–314. doi.org/10.22059/ijhst.2021.294838.331.
- Fadaei, M. and Akbarpour, V. (2023). Evaluation of the effect of ascorbic acid and sucrose foliar application on some quantitative and qualitative characteristics of cucurbita pepo var . *Styriaca*, 37(1): 89–104. doi.org/10.22067/jhs.2021.72192.1087.
- Fadam, M. A. N., & Hamad, M. R. (2023). Response of pomegranate seedlings cv.salimi to foliar spraying with chelated iron and nutrient solution prosol and proline. *Anbar Journal of Agricultural Sciences*, 21(2), 549-562. doi: 10.32649/ajas.2023.179754
- Fruhworth, G.O. and Hermetter, A. (2008). Production technology and characteristics of Styrian pumpkin seed oil, *European Journal of Lipid Science and Technology*, 110(7), pp. 637–644. doi.org/10.1002/ejlt.200700257.
- Habibi, A., Heidari, G., Sohrabi, Y. and Badakhshan, H. (2015). Effect of Different Fertilization Systems on Yield and Seed Mineral Elements of Pumpkin, *Iran Agricultural Research*, 33(2): 83–94.
- Hialy, M.Z.T. Al (2018). Influence of spraying with water extracts of liquorice and organic manures for animals on yield, and its components of trigonella foenum - graecum l. local variety. *Rafidain journal of science*, 28(3): 61-76.

- Hirzel, J., Millas, P., France, A. and Cerda, F. (2012). Compost tea effects on production and extraction of nitrogen in ryegrass cultivated on soil amended with commercial compost, *Compost Science and Utilization*, 20(2):97–104. doi.org/10.1080/1065657X.2012.10737032.
- Liu, X. Q., Xie, M. M., Hashem, A., Abd-Allah, E. F., & Wu, Q. S. (2023). Arbuscular mycorrhizal fungi and rhizobia synergistically promote root colonization, plant growth, and nitrogen acquisition. *Plant Growth Regulation*, 100(3), 691-701.
- Iswan, A. (2014). Agronomic traits and fruit quality of pineapple with different levels of chicken manure application, *Silpakorn U Science & Tech J*, 8(1): 67–73.
- Japelt, Rie Bak (2011). Vitamin D in plants Occurrence, analysis and biosynthesis. PhD thesis. *Division of Food Chemistry National Food Institute Technical University of Denmark*
- Jeznach, M., Danilcenko, H., Jariene, E., Kulaitiene, J. and Černiauskiene, J. (2012) Accumulation of antioxidative vitamins and minerals in seeds of oil pumpkin (*Cucurbita pepo* L. var. *styriaca*) cultivars, *Journal of Food, Agriculture and Environment*, 10(1):245–247.
- Khandaker, M.M., Azani, S., Majrashi, A., Alenazi, M.M., Saifuddin, M., Mohd, K.S. and Adnan, A.F.M. (2022). Morphophysiology, growth indices, and fruiting of pumpkin in response to organic amendments', *Semina: Ciencias Agrarias*, 43(3): 1211–1236. doi.org/10.5433/1679-0359.2022v43n3p1211.
- Kim, M. Y., Kim, E. J., Kim, Y. N., Choi, C., & Lee, B. H. (2012). Comparison of the chemical compositions and nutritive values of various pumpkin (Cucurbitaceae) species and parts. *Nutrition research and practice*, 6(1), 21.
- Kozhanova, L.A., G.A. Fedorova and g.i. Baram (2002). Determination of water- and fat-soluble vitamins in multivitamin preparations by high-performance liquid chromatography. *Journal of Analytical Chemistry*. 57(1): 40-45.
- Latifi, M.; A.B. Shahram and S.R. Lipayi (2012). Sowing date and plant population effects on seed yield of *Cucurbita pepo*. *International Journal of Agriculture and Biology*. 1814–9596.
- Najafi, S., Nasi, H.N., Tuncturk, R., Tuncturk, M., Sayyed, R.Z. and Amirnia, R. (2021) Biofertilizer application enhances drought stress tolerance and alters the antioxidant enzymes in medicinal pumpkin (*Cucurbita pepo* convar. *pepo* var. *styriaca*), *Horticulturae*, 7(12). doi.org/10.3390/horticulturae7120588.

- Oloyede, F.M.; O.C. Adebooye and E.M. Obuotor (2014). Planting date and fertilizer affect antioxidants in pumpkin fruit. *Scientia Horticulturae* 168: 46–50.
- Omidbeigi, R. (2004). Production and Process of Medicinal Herbs 3: 397. *Astane Ghods-e Razavi Publication*, Mashhad, Iran.
- Oyetunde, O., Benson, G. A. S., Oyetunde, A. K., Olalekan, O. J., Chukwu, C. U., & Anari, K. O. (2023). Agromorphological and nutritional quality profiles of fluted pumpkin (*Telfairia occidentalis* Hook F.) as influenced by cultivar, growing medium and soil amendment source. *International Journal of Horticultural Science*, 53-59.
- Payne R, Murray D, Harding S, Baird D, Soutou D (2009) GenStat for Windows 12th Edition Introduction: *VSN International, 5 The Waterhouse*. Waterhouse Street, Hemel Hempstead, Hertfordshire HP1 1ES, UK.
- Ros, M.; M. Hurtado-Navarro; A. Giménez; J.A. Fernández; C. Egea-Gilabert; P. Lozano-Pastor and J.A. Pascual (2020). Spraying agro-industrial compost tea on baby spinach crops: evaluation of yield, plant quality and soil health in field experiments. *Agronomy*, 10, 440: 1 – 15.
- Suhartini, S., Wijana, S., Surjono, Wardhani, N.W.S. and Muttaqin, S. (2020). Composting of chicken manure for biofertiliser production: A case study in Kidal Village, Malang Regency, *IOP Conference Series: Earth and Environmental Science*, 524(1). doi.org/10.1088/1755-1315/524/1/012016.