



IRAQI  
Academic Scientific Journals



العراقية  
المجلات الأكاديمية العلمية

TJAS

Tikrit Journal for  
Agricultural  
Sciences

ISSN:1813-1646 (Print); 2664-0597 (Online)

*Tikrit Journal for Agricultural Sciences*

Journal Homepage: <http://www.tjas.org>

E-mail: [tjas@tu.edu.iq](mailto:tjas@tu.edu.iq)

## The Role of Sumac Powder in Enhancing Physiological Parameters in Awassi Lambs

Mohammed S. Mohammed<sup>1</sup> , Mohhaiman A. Mohamad<sup>1</sup> , and Salih N. Hussein<sup>2</sup>

<sup>1</sup>Department of Animal Production, College of Agriculture, Tikrit University, Iraq

<sup>2</sup>Department of Publish Health, College of Veterinary, Tikrit University, Iraq

\*Correspondence email: [dr.mohsaleh@tu.edu.iq](mailto:dr.mohsaleh@tu.edu.iq)

### KEY WORDS:

Physiological; Sumac Powder;  
Lambs

Received: 27/03/2024

Revision: 26/07/2024

Proofreading: 22/09/2024

Accepted: 13/08/2024

Available online: 31/12/2024

©2024. This is an open access article under the CC by licenses

<http://creativecommons.org/licenses/by/4.0>



### ABSTRACT

The objective of the study was to know biological effect for the Sumac powder in the diets of Awassi lambs. A total of 24 Awassi lambs (12 male and 12 female) were used (age 3-4 months) ( $16.5 \pm 1.5$ g male and  $20 \pm 2.5$  g female) and divided to four groups (6 animals / 3 replicated). The experimental animals were fed into four individually in case ( $2.5 \times 1.5$  cm). Group 1 were fed on a diet without adding 0 % (control); groups 2, 3 and 4 were fed 1, 3 and 5% sumac powder. The results showed indicate that adding Sumac powder to the feed significant differences between the groups in both AST and ALT. There were not significant between the groups in creatinine, urea, total protein, albumin, and globulin levels in the blood serum of the animals compared to the control group. Our results suggested that the sumac could be used in the animal food without any effect on their health.

## دور مسحوق السماق في تحسين بعض الصفات الفسيولوجية في الحملان العواسية

محمد صالح محمد السلامي ، مهيمن عبد السلام محمد و صالح نجم حسين  
قسم الانتاج الحيواني ، كلية الزراعة ، جامعة تكريت ، العراق  
قسم الصحة العامة ، كلية الطب البيطري ، جامعة تكريت ، العراق

### INTROUCTION

*Rhus coriaria* plant, has garnered attention for its potential health benefits and applications in various fields, including nutrition and pharmacology (Jamir & Sibian, 2023). Sumac is increasingly used in food science, exemplified by its incorporation into gummy candies, which received high sensory ratings for taste and texture (Jamir & Sibian, 2023). Despite these benefits, some studies suggest that sumac may not significantly impact measures like body weight (Nouri et al., 2023). The chemical composition of sumac powder, reveals a rich profile of bioactive compounds and nutrients. This composition contributes to its culinary and medicinal applications, and the contains significant amounts of carbohydrates (approximately 66.05%), moisture (32.61%), and minimal levels of fat (0.52%) and protein (0.52%)(Jamir & Sibian, 2023), and it is a source of dietary fibers, which are essential for digestive health(Nagib, 2017).

Sumac is rich in phenolic compounds, including gallic acid and its derivatives, which are linked to various health benefits such as antioxidant and anti-inflammatory properties(Grassia et al., 2021; Khaled et al., 2023), flavonoids, particularly quercetin and myricetin, enhances its therapeutic potential (Grassia et al., 2021). Khaled et al (2023) a showed sumac exhibits strong antioxidant properties, making it beneficial for combating oxidative stress (Khaled et al., 2023). On another hands, some studies indicate that sumac powder can help reduce cholesterol levels, supporting cardiovascular health (Nagib, 2017). A study demonstrated that lambs receiving a diet supplemented with 3% sumac powder exhibited significant improvements in wool traits, including staple length and clean wool weight (Al-Saadi, 2023).

The combination of sumac and exogenous fibrolytic enzymes further amplified these benefits, highlighting its role as an effective appetite stimulant (Al-Saadi, 2023). Increasing levels of sumac powder in lamb diets led to improved digestibility of nutrients, with organic matter digestibility reaching up to 84.50% at 5% sumac inclusion (Hussein et al., 2022). Enhanced feed conversion ratios were also observed, indicating better nutrient utilization(Hussein et al.,

2022). Sumac supplementation positively influenced blood parameters, including hemoglobin and red blood cell counts (Hadi & Al-Saadi, 2024). While the benefits of sumac powder are evident, Hussein *et al.*, (2022) indicate that the effects on growth rates may not be as pronounced, suggesting that further research is needed to optimize its use in lamb nutrition. The aim of this study to role of the sumac as a medical plants added to the Awassi lambs diet and affected to the some physiological parameters.

**MATERIAL AND METHODS**

The current study's procedures were carried out in conformity with the guidelines of the Ethics Research Committee of the College of Agriculture, Tikrit University.

Twelve female Awassi lambs were used in this research, with ages ranging between (3-4) months approximately, and they were purchased from the local market. The research was conducted in the (Sumum) area of the Samarra city. The lambs were placed inside a large closed hall with a paved floor and divided into several pens of 4 \* 10 meters in size. One of them was exploited and inside it was placed cages measuring 2.5 \* 1.5 by 12 cages where the feeding was individual and each cage was equipped with a feeder. And a bowl to drink water. After purchasing the animals and transporting them to the fields, they were placed under observation for a period of ten days, in order to ensure that they were free of diseases, and were included in a health and preventive program for many common diseases in the region. Four rations were used in this research as shown in Table 1. The first ration included barley, wheat bran and soybean meal. In this diet, needs were taken into account.

**Table 1.** Experiment diet ingredients and percentage of use (A) Male (B) Female

Ingredients (%)	Treatments (A)				Treatments (B)			
	C-non	S <sub>1</sub>	S <sub>2</sub>	S <sub>4</sub>	S <sub>1</sub>	S <sub>2</sub>	S <sub>4</sub>	S <sub>1</sub>
Barley	40	40	40	39	40	40	40	40
Wheat bran	39	40	40	39	40	40	40	39
Soybean meal (SBM)	10	10	11	11	9	8	7	7
Corn	10	8	5	5	10	10	9	8
Sumac	0	1	3	5	0	1	3	5
Min. & Vit. Mixture	1	1	1	1	1	1	1	1

C-non: control; S<sub>1</sub>, S<sub>2</sub> and S<sub>3</sub>:Datie with 1,2 and 3 % sumac powder

## Hematology parameters

Blood samples (20 ml) were collected from the animals' jugular veins from all lambs, blood was collected into two different types of vacutainer collecting tubes: one type was coated with ethylene diamine tetra acetic acid (EDTA) as an anticoagulant for evaluating the hematological parameters, and the other type was plain vacutainer tubes for separating serum to measure the parameters, separate serum samples were kept at -20°C pending further examination. Total protein was determined calorimetrically using the Biuret method as described by Yousef *et al.*,(2006); Albumin was determined calorimetrically by the method described by Doumas *et al.*,(1971); Globulin was determined by subtracting albumin from total protein; ALT and AST were determined by White *et al.*,(1970). Blood urea concentration was estimated by Nessler's reaction (Tanis and Naylor, 1968).

## Statistical analysis

The SAS appropriate statistical program was used to do a one-way ANOVA on the data (SAS, 2005). The Duncan significant differences (Duncan, 1953) approach was used to find the variance in the mean ( $P \leq 0.05$ ).

## RESULTS AND DISSCUSION

The results of the statistical analysis in table (2) indicate that adding Sumac powder to the feed recorded significant differences between the groups in both AST and ALT. It is noted that the Sumac powder additions recorded an increase in the AST level compared to the control group. On the other hand, it is noted that the second group recorded the highest level of ALT compared to S<sub>1</sub> group, while no significant differences were recorded between it and the C-non (control) and S<sub>3</sub> group (table 2).

**Table 2:** Effect of the Sumac powder in the diets of ALT and AST in Awassi lambs

Groups	Parameters	
	AST (mg/dL)	ALT(mg/dL)
C-non	8.50 ± 1.44 b	34.50 ± 0.32 ab
S <sub>1</sub>	13.25 ± 4.97 a	38.00 ± 0.24 a
S <sub>2</sub>	13.00 ± 1.47 a	32.50 ± 0.49 b
S <sub>3</sub>	14.00 ± 2.79 a	34.25 ± 0.94 ab

C-non: control; S<sub>1</sub>, S<sub>2</sub> and S<sub>3</sub>:Datie with 1,2 and 3 % sumac powder

The results of the statistical analysis in table (3) indicate that adding Sumac powder to the feed recorded non-significant differences between the groups in creatinine and urea levels in the blood serum of the animals compared to the control group.

**Table 3:** Effect of the Sumac powder in the diets of Creatinine and Urea in Awassi lambs

Groups	Parameters	
	Creatinine (mg/dL)	Urea (mg/dL)
C-non	0.99 ± 0.03 a	41.65 ± 5.92 a
S <sub>1</sub>	1.09 ± 0.10 a	52.78 ± 7.16 a
S <sub>2</sub>	1.07 ± 0.04 a	43.64 ± 4.70 a
S <sub>3</sub>	0.97 ± 0.03 a	51.33 ± 4.43 a

C-non: control; S<sub>1</sub>, S<sub>2</sub> and S<sub>3</sub>:Datie with 1,2 and 3 % sumac powder

On another hands, the results in table (4) showed non-significant differences between the groups in total protein, albumin, and globulin levels in the blood serum of the animals compared to the control group.

**Table 4:** Effect of the Sumac powder in the diets of total protein, albumin, globulin, and ALB/GLU in Awassi lambs

Groups	Parameters			
	Total Protein (g/dL)	Albumin (g/dL)	Globulin (g/dL)	ALB/GLU
C-non	6.59 ± 0.13	3.12 ± 0.31	3.47 ± 0.32	0.95 ± 0.21
S <sub>1</sub>	6.56 ± 0.11	2.21 ± 0.31	4.35 ± 0.24	0.52 ± 0.10
S <sub>2</sub>	7.19 ± 0.22	2.55 ± 0.59	4.63 ± 0.49	0.61 ± 0.18
S <sub>3</sub>	6.80 ± 0.27	2.99 ± 0.65	3.80 ± 0.90	0.99 ± 0.31

C-non: control; S<sub>1</sub>, S<sub>2</sub> and S<sub>3</sub>:Datie with 1,2 and 3 % sumac powder

Our results showed that sumac powder effect to the liver enzyme, and one of the primary factors may be the antioxidant properties of Sumac, which is rich in phenolic compounds and flavonoids known to mitigate oxidative stress in the liver (Chung *et al.*, 2012). Reducing oxidative stress can improve liver function, which may lead to changes in enzyme levels (Liu *et al.*, 2018). Sumac’s antioxidant properties are impressive, but its anti-inflammatory effects are also important. Research suggests that the bioactive compounds in sumac can help reduce inflammation, a condition often associated with high liver enzyme levels (Muhaimed *et al.*, 2021). By reducing inflammation in the liver, sumac may positively impact liver enzyme activity, supporting its role in promoting liver health. In addition, the nutritional composition of sumac is important. A variety of bioactive compounds can influence metabolic processes in animals, leading to changes in enzyme activity (Khan *et al.*, 2018). Dietary changes can significantly impact nutrient absorption and metabolism, which may indirectly impact liver enzyme levels (Hajialyani *et al.*, 2019). Furthermore, sumac may help promote healthy gut microbiota, which is crucial for healthy liver function through gut-liver communication. A balanced gut microbiome is associated with improved metabolism and overall liver health, which may explain the observed changes in AST

and ALT levels (Lau et al., 2021). One possible reason for the results in this study is that the concentrations of sumac powder used (1%, 3%, and 5%) may not have been high enough to significantly impact protein synthesis or metabolism. Previous studies suggest that dietary changes often need to exceed a certain level to meaningfully impact protein levels (Tieland et al., 2017). Additionally, the proteins in the animals' diets may have been sufficient to meet their physiological needs, which may have masked any effects from sumac. It is also worth noting that sumac is primarily known for its antioxidant and anti-inflammatory benefits rather than its effects on protein metabolism. While its bioactive components may support liver health, they may not directly impact pathways involved in the synthesis or degradation of serum proteins (Khan et al., 2020; Kamil et al., 2021). This means that while sumac may enhance liver function, it does not necessarily lead to changes in total protein levels or its various fractions in the bloodstream. Furthermore, the physiological state of the animals could also be a factor. Since the lambs in this study were young and relatively healthy, their protein metabolism may have been stable and less affected by feed additives. Factors such as age and health status can significantly influence protein metabolism and serum protein synthesis (Hajjalayani et al., 2019; Hussein et al., 2022). Finally, it is important to consider that the balance of different nutritional components can affect protein levels. The overall composition of the diet, including other feed components, may have mitigated any potential effects of sumac on protein metabolism (Portugal et al., 2021).

## **CONCLUSION**

This study revealing significant findings regarding on liver enzymes, no significant changes were observed in other important blood parameters compared to the control group. Sumac powder shows promise as a beneficial dietary supplement in lamb nutrition, warranting further investigation into its long-term effects and potential benefits

## **CONFLICT OF INTEREST**

The authors declare no conflicts of interest associated with this manuscript.

## ACKNOWLEDGMENTS

The authors gratefully acknowledge the staff of the Department of Animal Production, College of Agriculture, Tikrit University for their technical and general support.

## REFERENCES

- Tieland, M., Franssen, R., Dullemeijer, C., van Dronkelaar, C., Kim, H. K., Ispoglou, T., ... & de Groot, L. C. (2017). The impact of dietary protein or amino acid supplementation on muscle mass and strength in elderly people: individual participant data and meta-analysis of RCT's. *The Journal of nutrition, health and aging*, 21(9), 994-1001.
- Lau, A. W. Y., Tan, L. T. H., Ab Mutalib, N. S., Wong, S. H., Letchumanan, V., & Lee, L. H. (2021). The chemistry of gut microbiome in health and diseases. *Progress In Microbes & Molecular Biology*, 4(1).
- Chung, K. T., Wong, T. Y., & Wei, C. I. (2012). Tannins and human health: A review. *Critical Reviews in Food Science and Nutrition*, 52(3), 201-213.
- Doumas, B.T.; Ard Watson, W.; Biggs, H.G. Albumin standards and the measurement of serum albumin with bromcresol green. *Clin. Chim. Acta* 1971, 31, 87–96, doi:10.1016/0009-8981(71)90365-2.
- Gülderen, Yilmaz., Gülnur, Ekşi., Betül, Demirci., Fatih, Demirci. (2020). 4. Chemical characterization of the fatty acid compositions and antimicrobial activity of sumac (*rhus coriaria* l.) fruits, growing naturally in turkey and sold in herbalist markets. doi: 10.33483/JFPAU.645467
- Hajialyani, M., Hosein Farzaei, M., Echeverría, J., Nabavi, S., Uriarte, E. and Sobarzo-Sánchez, E., 2019. Hesperidin as a neuroprotective agent: a review of animal and clinical evidence. *Molecules* 24(3): 648. <https://doi.org/10.3390/molecules24030648>
- Ibrahim, M., Abu-Reida., Rana, M., Jamous., Mohammed, S., Ali-Shtayeh. (2014). 7. Phytochemistry , Pharmacological Properties and Industrial Applications of *Rhus Coriaria* L. ( Sumac ). *jordan journal of biological sciences*, doi: 10.12816/0008245

- Kamil, Y. M., Katab, T., & Mohammed, A. B. M. (2021, May). Effect of Supplementing laying Japanese quail (*Coturnix coturnix japonica*) diets with Amla and Green Tea Extracts on the Product Performance and Biochemical Parameters. In *IOP Conference Series: Earth and Environmental Science* (Vol. 761, No. 1, p. 012109). IOP Publishing.
- Katia, Arena., Emanuela, Trovato., Francesco, Cacciola., Ludovica, Spagnuolo., Elisa, Pannucci., P., Guarnaccia., Luca, Santi., Paola, Dugo., Luigi, Mondello., Laura, Dugo. (2022). 2. Phytochemical Characterization of *Rhus coriaria* L. Extracts by Headspace Solid-Phase Micro Extraction Gas Chromatography, Comprehensive Two-Dimensional Liquid Chromatography, and Antioxidant Activity Evaluation. *Molecules*, doi: 10.3390/molecules27051727
- Khan, M., Ahmad, K., Hassan, S., Imran, M., Ahmad, N. and Xu, C., 2018. Effect of novel technologies on polyphenols during food processing. *Innovative Food Science and Emerging Technologies* 45: 361–381. <https://doi.org/10.1016/j.ifset.2017.12.006>
- Khan, M. I., et al. (2020). Nutritional and health benefits of Sumac (*Rhus coriaria*): A review. *Journal of Food Science and Technology*, 57(1), 1-10.
- Liu, W., et al. (2018). Antioxidant and anti-inflammatory effects of natural products in the prevention of liver diseases: A review. *Food Chemistry*, 265, 214-222.
- Mohammad, Mahdi, Ahmadian-Attari., Mohsen, Amini., Hassan, Farsam., Gholamreza, Amin., Mohammad, Reza, Fazeli., Hamid, Reza, Monsef, Esfahani., Hossein, Jamalifar., Amir, Bairami. (2016). 8. Isolation of Major Active Antibacterial Compounds of Sumac Fruit (*Rhus coriaria* L.). *International Journal of Enteric Pathogens*, doi: 10.15171/IJEP.2016.11
- Musa, Özcan., Haydar, Hacıseferoğulları. (2004). 3. A condiment (sumac (*rhus coriaria* l.) fruits): some physico-chemical properties.
- Nagib, M. (2017). 1. Hypolipidemic Effect of Sumac (*Rhus Coriaria* L) Fruit Powder and Extract on Rats Fed High Cholesterol Diet. doi: 10.21608/BNNI.2017.6726
- Parisa, Moghadam., Shahram, Dadelahi., Yasamin, Sayed, Hajizadeh., Milad, G., Matin., Milad, Amini., Saba, Hajazimian. (2020). 6. Chemical Composition and Antibacterial Activities



of Sumac Fruit (*Rhus coriaria*) Essential Oil on Dental Caries Pathogens. The Open Microbiology Journal, doi: 10.2174/1874285802014010142.

Portugal, T. B., Szymczak, L. S., de Moraes, A., Fonseca, L., Mezzalira, J. C., Savian, J. V., ... & Monteiro, A. L. G. (2021). Low-Intensity, high-frequency grazing strategy increases herbage production and beef cattle performance on sorghum pastures. *Animals*, 12(1), 13.

White, B.W.; Saunders, F.A.; Scadden, L.; Bach-Y-Rita, P.; Collins, C.C. Seeing with the skin. *Percept. Psychophys.* 1970, 7, 23–27, doi:10.3758/BF03210126.

Yousef, M.I.; El-Demerdash, F.M. Acrylamide-induced oxidative stress and biochemical perturbations in rats. *Toxicology* 2006, 219, 133–141, doi:10.1016/j.tox.2005.11.008.

Muhaimeed, A. R., Hussein, S. N., Ibraheem, M. W., & Muhammed, A. M. (2021, April). Effect of Different Levels of *Rhus Coriaria* L. powder in Awassi Lambs Diet on Some Carcass Characteristics. In *IOP Conference Series: Earth and Environmental Science* (Vol. 735, No. 1, p. 012013). IOP Publishing.

Hussein, S. N., Mohammad, M. A. A., Aljumaily, M. H., Mohammed, M. S., & Abass, K. S. (2022, July). Effect of Different Levels of Red Sumac Powder *Rhus Coriria* L. on Productive Performance, Coefficient of Digestion And Some Rumen Fluid Characteristics of Awassi Lambs Females. In *IOP Conference Series: Earth and Environmental Science* (Vol. 1060, No. 1, p. 012073). IOP Publishing.